

# **THE INFLUENCE OF EVALUATION PROTOCOL ON TIME SPENT EXERCISING AT A HIGH LEVEL OF OXYGEN UPTAKE DURING CONTINUOUS CYCLING**

K.L. Merry<sup>1</sup>, M. Glaister<sup>2</sup>, G. Howatson<sup>3,4</sup>, K. van Someren<sup>5</sup>

1. School of Science, University of Derby, Derby, United Kingdom
2. School of Sport, Health and Applied Science, St Mary's University College, Twickenham, United Kingdom
3. School of Life Sciences, Northumbria University, Newcastle upon Tyne, United Kingdom
4. Centre for Aquatic Research, University of Johannesburg, Johannesburg, South Africa
5. GlaxoSmithKline Human Performance Laboratory, Brentford , United Kingdom

Congresses: None

Funding: None

Conflicts of interest: None

Acknowledgments: None

Corresponding Author:

K.L. Merry

School of Science

University of Derby,

Kedleston Road,

DE22 1GB, Derby

United Kingdom,

k.merry@derby.ac.uk;

## ABSTRACT

**Aim:** This study evaluated the effects of protocol variation on the time spent exercising at  $\geq 95\%$   $\text{VO}_{2\text{max}}$  during cycle ergometer trials performed at the exercise intensity associated with  $\text{VO}_{2\text{max}}$  ( $\text{iVO}_{2\text{max}}$ ). **Methods:** Nine male triathletes (age:  $32 \pm 10$  years; body mass:  $73.3 \pm 6.1$  kg; stature:  $1.79 \pm 0.07$  m;  $\text{VO}_{2\text{max}}$ :  $3.58 \pm 0.45$  L.min<sup>-1</sup>) performed four exercise tests. During tests 1 and 2, participants performed a maximal incremental cycle ergometer test using different stage durations (1 min and 3 min) for the determination of  $\text{iVO}_{2\text{max}(1 \text{ min})}$  and  $\text{iVO}_{2\text{max}(3 \text{ min})}$ . During tests 3 and 4, participants performed a continuous bout of exhaustive cycling at  $\text{iVO}_{2\text{max}(1 \text{ min})}$  (CONT<sub>1</sub>) and  $\text{iVO}_{2\text{max}(3 \text{ min})}$  (CONT<sub>3</sub>). **Results:**  $\text{iVO}_{2\text{max}(1 \text{ min})}$  was significantly greater ( $P < 0.001$ ) than  $\text{iVO}_{2\text{max}(3 \text{ min})}$  ( $340 \pm 31$  W vs.  $299 \pm 44$  W). Time to exhaustion (TTE) measured during CONT<sub>3</sub> was significantly longer ( $P < 0.001$ ) than CONT<sub>1</sub> ( $529 \pm 140$  s vs.  $214 \pm 65$  s). Time spent at  $\text{VO}_{2\text{max}}$  was significantly longer ( $P = 0.036$ ) during CONT<sub>3</sub> than CONT<sub>1</sub> ( $146 \pm 158$  s vs.  $11 \pm 20$  s), and time spent at  $\geq 95\%$   $\text{VO}_{2\text{max}}$  was significantly longer ( $P = 0.005$ ) during CONT<sub>3</sub> than CONT<sub>1</sub> ( $326 \pm 211$  s vs.  $57 \pm 51$  s). **Conclusion:** These results show that when exercising continuously at  $\text{iVO}_{2\text{max}}$ , time spent at  $\geq 95\%$   $\text{VO}_{2\text{max}}$  is influenced by the initial measurement of  $\text{iVO}_{2\text{max}}$ .

Keywords: Continuous exercise, maximal oxygen uptake, time spent at maximal oxygen uptake, time to exhaustion

## Introduction

Several researchers have suggested that it may be necessary for highly trained athletes to elicit  $\text{VO}_{2\text{max}}$  during training sessions in order to improve it.<sup>1-4</sup> Consequently, an interest has been developed in determining training protocols that allow athletes to spend the longest time

exercising at  $\text{VO}_{2\text{max}}$  ( $\text{T@VO}_{2\text{max}}$ ) or near to ( $\geq 95\%$ )  $\text{VO}_{2\text{max}}$  ( $\text{T@95\%VO}_{2\text{max}}$ ) during continuous and intermittent exercise.

The exercise intensity at  $\text{VO}_{2\text{max}}$  ( $\text{iVO}_{2\text{max}}$ ) has been defined in several ways by various authors but generally represents the minimum exercise intensity associated with the attainment of  $\text{VO}_{2\text{max}}$  during a maximal incremental test.<sup>5 6</sup> Measurement of  $\text{iVO}_{2\text{max}}$  in this way has proved popular with coaches and researchers when setting interval training intensities for athletes.<sup>7-9</sup>

The exercise intensity at  $\text{VO}_{2\text{max}}$  may allow the longest time to be spent exercising at or near to  $\text{VO}_{2\text{max}}$  during continuous exercise<sup>10</sup> and several studies have reported that  $\text{iVO}_{2\text{max}}$  and intensities very close to  $\text{iVO}_{2\text{max}}$  (95 – 105%  $\text{iVO}_{2\text{max}}$ ) allow the longest time to be spent exercising at or near to  $\text{VO}_{2\text{max}}$  during intermittent exercise.<sup>11-13</sup> However, the protocol to measure  $\text{iVO}_{2\text{max}}$  has differed between studies. Some studies have measured  $\text{iVO}_{2\text{max}}$  using continuous incremental tests with stage durations ranging from 1 to 3 min<sup>9 13-20</sup> whilst others have measured  $\text{iVO}_{2\text{max}}$  using discontinuous protocols with stage durations ranging from 2 – 3 min.<sup>11 21-23</sup> The disparity between studies in terms of the method of  $\text{iVO}_{2\text{max}}$  determination makes comparison of  $\text{iVO}_{2\text{max}}$  values and indeed  $\text{T@VO}_{2\text{max}}$  difficult. It has been demonstrated that an  $\text{iVO}_{2\text{max}}$  test protocol including 1 min stage durations did not produce significantly different values for  $\text{iVO}_{2\text{max}}$  or the exercise time to exhaustion (TTE) at  $\text{iVO}_{2\text{max}}$  than a separate protocol with 2 min stage durations. However,  $\text{T@VO}_{2\text{max}}$  was not measured in the study.<sup>15</sup> It is clear that different protocols for  $\text{iVO}_{2\text{max}}$  determination can produce different  $\text{iVO}_{2\text{max}}$  values. A previous study compared a continuous incremental test protocol with stages of 1 min, with two discontinuous incremental test protocols with stages of 2 and 3 min respectively.<sup>23</sup> Significant differences in  $\text{iVO}_{2\text{max}}$  were reported between each of the three protocols. However,  $\text{T@VO}_{2\text{max}}$  was also not measured.

It has been recommended that for running-based assessment,  $iVO_{2max}$  should be measured during a continuous incremental test with stage durations of one minute.<sup>15</sup> However, some researchers have argued that maximal incremental tests should utilise stages of at least 3 minutes to obtain a stabilisation of gas exchange parameters and an optimal relationship between  $VO_2$  and exercise intensity.<sup>5 24-26</sup> If insufficient time is given for the elicitation of  $VO_{2max}$  during a given stage at the end of an incremental test, then  $iVO_{2max}$  may be reported as the intensity of the following stage and subsequently overestimated. Accuracy in determining  $iVO_{2max}$  is important, since an overestimation of  $iVO_{2max}$  may influence the total time spent exercising at  $VO_{2max}$ , compromising potential training adaptations. With this in mind, there is a need to clarify the impact of  $iVO_{2max}$  test protocol differences on  $T@VO_{2max}$ , since no common protocol for  $iVO_{2max}$  measurement currently exists. Therefore, the aim of this study is to assess the length of time spent exercising at a high level of  $VO_2$  ( $\geq 95\% VO_{2max}$ ) during continuous cycling to exhaustion at  $iVO_{2max}$  determined using two different protocols.

## **Materials and Methods**

### *Subjects*

Subjects were nine male triathletes. All had been training and competing in triathlon events for at least 3 years and had been engaged in  $\geq 4$  hours of training per week including running, cycling and swimming during the 6 months prior to testing. The subjects possessed the following characteristics (values are expressed as mean  $\pm$  standard deviation); age:  $32 \pm 10$  years, body mass:  $73.3 \pm 6.1$  kg, stature:  $1.79 \pm 0.07$  m,  $VO_{2max}$ :  $3.58 \pm 0.45$  L.min<sup>-1</sup>. All experimental procedures and potential discomforts associated with the study were explained to subjects before they gave written informed consent to participate. The study was approved by Loughborough College's Research Ethics Committee.

### *Experimental overview*

Each subject reported to the laboratory on five separate occasions at approximately the same time of day. Each visit was separated by at least 48 hours and all visits were completed within two weeks. During visit one, participants performed a submaximal incremental test for the determination of the onset of blood lactate accumulation (OBLA). During each of visits two and three, a single maximal incremental test was performed to determine  $\text{VO}_{2\text{max}}$  and  $\text{iVO}_{2\text{max}}$ . Each of the maximal incremental tests incorporated different stage durations (1 and 3 min) and were performed in a randomised, counterbalanced order for the determination of  $\text{iVO}_{2\text{max}(1 \text{ min})}$  and  $\text{iVO}_{2\text{max}(3 \text{ min})}$ . During visits four and five, participants performed two continuous cycle exercise to exhaustion tests at  $\text{iVO}_{2\text{max}(1 \text{ min})}$  ( $\text{CONT}_1$ ) and  $\text{iVO}_{2\text{max}(3 \text{ min})}$  ( $\text{CONT}_3$ ) for the determination of  $\text{T@VO}_{2\text{max}}$ . The  $\text{CONT}_1$  and  $\text{CONT}_3$  tests were performed in a randomised, counterbalanced order.

### *Procedures*

All testing was performed on an electro-magnetically braked cycle ergometer (Lode Excalibur Sport, Groningen, The Netherlands) modified with cleat pedals and low-profile racing handlebars. The riding position, including saddle and handlebar positions was adjusted to resemble each subjects own bike and was replicated in subsequent trials. During all tests respiratory gas exchange was measured at the mouth using a breath by breath online gas analysis system (Cortex Metalyzer 3B, Leipzig, Germany). This system has been shown to have very good repeatability for measures of  $\text{VO}_2$  ( $\text{ICC} = 0.964$ ).<sup>27</sup> During measurement, subjects wore a rubber face mask (Hans Rudolph 8940, Hans Rudolph, Inc., Kansas City, USA) and breathed through a low resistance volume transducer (Triple V, Erich Jaeger GmbH, Hoechberg, Germany). Immediately before each test, the gas analyser was calibrated using

ambient air which was assumed to contain 20.93% O<sub>2</sub> and 0.03% CO<sub>2</sub>, and with a gas of known O<sub>2</sub> and CO<sub>2</sub> concentrations. Heart rate (HR) was recorded continuously throughout each test using a telemetric system which was interfaced with the gas analyser (Polar Electro Oy, Kempele, Finland). Ear lobe capillary blood samples were obtained in a 30 s pause between stages during the submaximal incremental test, and immediately after each subsequent maximal incremental test and CONT<sub>1</sub> and CONT<sub>3</sub> tests and analysed for lactate using an automated analyser (Biosen 5030, EKF Industrie, Germany). The repeatability of this analyser has previously been reported ( $R^2 = 0.99$ ; CV = 1.4%).<sup>28</sup>

#### *Submaximal Incremental Test*

The submaximal incremental test commenced at a workload equal to 2 W.kg bm<sup>-1</sup> and consisted of work rate increments of 20 W every 3 min. The test ended following attainment of OBLA which was considered to be equal to a blood lactate concentration of 4 mmol.L<sup>-1</sup>.<sup>29</sup> Blood lactate was plotted against power output and the power output that corresponded to OBLA was identified using specific software.<sup>30</sup>

#### *Determination of iVO<sub>2max</sub>*

For iVO<sub>2max(1 min)</sub> and iVO<sub>2max(3 min)</sub>, testing commenced at a workload 80 W less than the workload associated with OBLA. Power output was increased by 20 W every 1 min thereafter until volitional fatigue during iVO<sub>2max(1 min)</sub> and 20 W every 3 min for iVO<sub>2max(3 min)</sub>. iVO<sub>2max(1 min)</sub> and iVO<sub>2max(3 min)</sub> were taken as the lowest exercise intensities that elicited VO<sub>2max</sub> as long as this intensity was held for 60 s. If VO<sub>2max</sub> was elicited during a stage that wasn't maintained for 60 s then the intensity of the preceding stage was taken as iVO<sub>2max</sub> in line with previous recommendations.<sup>15</sup> During all tests, subjects remained seated and cycled at their preferred cycling cadence which was held constant for all visits. For all subjects this ranged from 85 – 90

rpm. Testing was terminated if cycling cadence changed by more than 10 rpm.  $\text{VO}_{2\text{max}}$  was defined by the following criteria: 1) oxygen consumption stopped increasing linearly with exercise; 2) Heart rate reached within  $10 \text{ beats} \cdot \text{min}^{-1}$  of age predicted maximum heart rate; 3) Respiratory exchange ratio (RER) was  $\geq 1.15$ ; 4) blood lactate concentration upon completion of the test was  $\geq 8 \text{ mmol} \cdot \text{L}^{-1}$  (34). Following data collection, all  $\text{VO}_2$  values were filtered and smoothed using rolling 30 s averages.  $\text{VO}_{2\text{max}}$  was taken as the highest 30 s  $\text{VO}_2$  value obtained during each test.

#### *Determination of Time at $\text{VO}_{2\text{max}}$*

Immediately before the  $\text{CONT}_1$  and  $\text{CONT}_3$  tests, participants completed a standardised warm up consisting of 10 min of cycling at 150 W. Prior to the start of each test, the exercise intensity corresponding to  $\text{iVO}_{2\text{max}}(1 \text{ min})$  or  $\text{iVO}_{2\text{max}}(3 \text{ min})$  was manually imposed onto the flywheel of the cycle ergometer using the ergometer's external control unit. At the start of each  $\text{CONT}_1$  and  $\text{CONT}_3$  test, participants were encouraged to obtain their preferred cycling cadence as quickly as possible. Timing began when the ergometer's external control unit displayed the appropriate power output value corresponding to  $\text{iVO}_{2\text{max}}(1 \text{ min})$  or  $\text{iVO}_{2\text{max}}(3 \text{ min})$  (for all participants this occurred within 15 s). Participants were given strong verbal encouragement throughout the test to induce a maximal effort. Time to exhaustion (TTE) during each test was recorded.  $\text{T@VO}_{2\text{max}}$  and  $\text{T@95\%VO}_{2\text{max}}$  were calculated through the accumulation of  $\text{VO}_2$  values superior or equal to 100% and 95% of the mean  $\text{VO}_{2\text{max}}$  calculated from the maximal incremental tests. Time to achieve  $\text{VO}_{2\text{max}}$  ( $\text{TTAVO}_{2\text{max}}$ ) and time to achieve 95%  $\text{VO}_{2\text{max}}$  ( $\text{TTA95\%VO}_{2\text{max}}$ ) were defined as the time taken to achieve 100% and 95% of the mean  $\text{VO}_{2\text{max}}$  calculated from the maximal incremental tests.

#### *Statistical Analysis*

Statistical analyses were performed using the Statistical Package for the Social Sciences (SPSS for Windows, SPSS Inc., Chicago, IL). Prior to statistical analysis, all measures were assessed for normality using the Shapiro-Wilk (W) test.<sup>32</sup> Mean values for  $iVO_{2max(1\text{ min})}$  and  $iVO_{2max(3\text{ min})}$  were compared using a paired samples t-test. Values for  $VO_{2max}$  measured during  $iVO_{2max(1\text{ min})}$  and  $iVO_{2max(3\text{ min})}$  were also compared using a paired samples t-test. Since it was found that the data for  $T@VO_{2max}$  was not normally distributed, mean values for  $T@VO_{2max}$  measured during  $CONT_1$  and  $CONT_3$  were compared using a Wilcoxon matched pairs test. Mean values for TTE,  $T@95\%VO_{2max}$ ,  $TTAVO_{2max}$ ,  $TTA\%95VO_{2max}$  and blood lactate concentration (BLa) measured during  $CONT_1$  and  $CONT_3$  were compared using paired samples t-tests. Mean values for  $TTAVO_{2max}$  and  $TTA95\%VO_{2max}$  expressed as a percentage of TTE were compared using paired t-tests. The above analyses provided 95% confidence limits for all estimates. For all analyses, results were considered significant at  $P < 0.05$ . All results and descriptive statistics are reported as means  $\pm$  standard deviations.

## Results

### *Maximal incremental tests*

Mean values for  $iVO_{2max(1\text{ min})}$ ,  $iVO_{2max(3\text{ min})}$  and their corresponding  $VO_{2max}$  values are summarised in Table 1.  $iVO_{2max}$  was significantly greater ( $P < 0.001$ ) (mean difference 41 W; 95% likely range: 29 – 54 W) during  $iVO_{2max(1\text{ min})}$  than  $iVO_{2max(3\text{ min})}$ . Mean  $VO_{2max}$  values obtained during the incremental tests were not significantly different ( $P > 0.05$ ).

### *Continuous cycling to exhaustion*

The mean values for each of the parameters measured during  $CONT_1$  and  $CONT_3$  tests are presented in Table 2. The mean TTE measured during  $CONT_3$  was significantly longer ( $P < 0.001$ ) (mean difference 315 s; 95% likely range: 209 – 424 s) than the TTE measured during  $CONT_1$ . Six participants elicited a  $VO_2 \geq VO_{2max}$  during  $CONT_1$ . Seven participants elicited a



$\text{VO}_2 \geq \text{VO}_{2\text{max}}$  during  $\text{CONT}_3$ .  $\text{T@VO}_{2\text{max}}$  measured during  $\text{CONT}_3$  was significantly longer ( $P = 0.036$ ) (mean difference 134 s; 95% likely range: 8 – 260 s) than  $\text{T@VO}_{2\text{max}}$  measured during  $\text{CONT}_1$ . Six participants elicited a  $\text{VO}_2 \geq 95\%$  of  $\text{VO}_{2\text{max}}$  during  $\text{CONT}_1$ . All participants elicited a  $\text{VO}_2 \geq 95\%$  of  $\text{VO}_{2\text{max}}$  during  $\text{CONT}_3$ . Subsequently,  $\text{T@95\%VO}_{2\text{max}}$  was significantly longer ( $P = 0.005$ ) (mean difference 269 s; 95% likely range: 110 – 428 s) during  $\text{CONT}_3$  than  $\text{CONT}_1$ .  $\text{TTAVO}_{2\text{max}}$  and  $\text{TTA95\%VO}_{2\text{max}}$  were both significantly shorter ( $P < 0.001$  and  $P = 0.029$ ) (mean difference 222 s; 95% likely range: 144 – 299 s) (mean difference 76 s; 95% likely range: 10 – 142 s) during  $\text{CONT}_1$  than during  $\text{CONT}_3$ . Mean BLa measured upon termination of  $\text{CONT}_1$  was significantly greater than the BLa measured upon termination of  $\text{CONT}_3$  ( $P < 0.001$ ) (mean difference 1.8 mMol.L<sup>-1</sup>; 95% likely range: 1.0 – 2.5 mMol.L<sup>-1</sup>). When expressed as a percentage of TTE,  $\text{TTAVO}_{2\text{max}}$  ( $91 \pm 27$  vs.  $77 \pm 17\%$ ) and  $\text{TTA95\%VO}_{2\text{max}}$  ( $46 \pm 11$  vs.  $34 \pm 16\%$ ) were not significantly different ( $P > 0.05$ ) between  $\text{CONT}_1$  and  $\text{CONT}_3$ .

## Discussion

The purpose of this study was to assess the influence of  $\text{iVO}_{2\text{max}}$  test protocol differences on the time spent exercising at a high percentage of  $\text{VO}_{2\text{max}}$  during continuous cycling to exhaustion at  $\text{iVO}_{2\text{max}}$ . The main finding was that a significantly longer time was spent exercising at a high percentage of  $\text{VO}_{2\text{max}}$  when  $\text{iVO}_{2\text{max}}$  was determined using a protocol involving stage durations of 3 min, compared to a protocol involving stage durations of 1 min.

As expected,  $\text{iVO}_{2\text{max}(1 \text{ min})}$  represented a significantly greater exercise intensity than  $\text{iVO}_{2\text{max}(3 \text{ min})}$ . This is consistent with other studies that have measured  $\text{iVO}_{2\text{max}}$  using several protocols with different stage durations.<sup>23</sup> Although  $\text{iVO}_{2\text{max}(1 \text{ min})}$  and  $\text{iVO}_{2\text{max}(3 \text{ min})}$  could both be characterised as 100%  $\text{iVO}_{2\text{max}}$ , it is clear that  $\text{iVO}_{2\text{max}(1 \text{ min})}$  is not the minimal exercise intensity associated with  $\text{VO}_{2\text{max}}$  since it produced intensities 12% greater than  $\text{iVO}_{2\text{max}(3 \text{ min})}$ .

The mean TTE when cycling continuously at  $iVO_{2max(3\text{ min})}$  (CONT<sub>3</sub>) was over twice as long as that recorded during the continuous cycling to exhaustion at  $iVO_{2max(1\text{ min})}$  (CONT<sub>1</sub>) and reflects the substantial effect that small differences in  $iVO_{2max}$  determinations can have on TTE at  $iVO_{2max}$  during continuous exercise. Previous research has demonstrated an inverse relationship between  $iVO_{2max}$  and TTE in homogenous samples of athletes.<sup>6 14 33</sup> Since  $iVO_{2max(1\text{ min})}$  represented a higher exercise intensity than  $iVO_{2max(3\text{ min})}$ , it is anticipated that the depletion of the anaerobic energy reserves would have been greater during CONT<sub>1</sub> than during CONT<sub>3</sub> increasing the rate at which metabolites associated with fatigue ( $P_i$ ,  $H^+$ ,  $K^+$ ) accumulate.<sup>34 35</sup> The significantly greater BLa values recorded following CONT<sub>1</sub> support this premise.

TTE values reported for CONT<sub>3</sub> are higher than mean values previously reported for elite cyclists (222 – 245 s),<sup>14 36</sup> triathletes (390 s),<sup>37</sup> and competitive cyclists (224 - 239 s).<sup>38</sup> Since the subjects used in the present study can be considered less well trained than those used in previous studies, the longer TTE values reported for CONT<sub>3</sub> are likely due to differences in the protocols used to determine  $iVO_{2max}$ . Previous studies have employed shorter stage durations (15 – 120 s),<sup>14 36 38</sup> larger increments (35 – 50 W)<sup>14 37</sup> or a combination of shorter stage durations and larger increments<sup>14</sup> than those used during  $iVO_{2max(3\text{ min})}$ . Therefore, it is likely that  $iVO_{2max(3\text{ min})}$  represents a lower relative exercise intensity than the  $iVO_{2max}$  utilised in other studies and likely accounts for the longer TTE values.

The mean  $TTAVO_{2max}$  during CONT<sub>3</sub> was significantly longer than that recorded during CONT<sub>1</sub>. The accelerated  $VO_2$  kinetics during CONT<sub>1</sub> is consistent with the higher exercise intensity utilised during CONT<sub>1</sub>. Several studies that have compared continuous exercise at intensities  $\leq iVO_{2max}$  with continuous exercise at intensities  $> iVO_{2max}$  have reported faster

TTAVO<sub>2max</sub> values for intensities > iVO<sub>2max</sub>.<sup>33 39</sup> Hence, it was expected that TTAVO<sub>2max</sub> would be faster during exercise at iVO<sub>2max</sub>(1 min) than during exercise at iVO<sub>2max</sub>(3 min). Oxidative energy production and the speed of the rise in oxidative metabolism at the onset of exercise appear to be increased as exercise intensity is increased.<sup>40 41</sup> Oxidative phosphorylation may be stimulated by an increase in the intracellular concentrations of ADP, Pi and Ca<sup>2+</sup>.<sup>40 42 43</sup> Since the rate of ATP hydrolysis and its subsequent resynthesis from PCr is proportional to exercise intensity, it is hypothesised that exercise during CONT<sub>1</sub> may have elevated those stimuli responsible for an increase in oxidative phosphorylation in comparison to CONT<sub>3</sub>.

The fact that T@VO<sub>2max</sub> and T@95% VO<sub>2max</sub> were both significantly longer during CONT<sub>3</sub> than CONT<sub>1</sub> likely reflect the longer TTE values recorded during CONT<sub>3</sub> compared to CONT<sub>1</sub> and are supported by previous research which has reported significant correlations between TTE and T@VO<sub>2max</sub> ( $r = 0.89$ ,  $P < 0.05$ ).<sup>19</sup> Furthermore, several studies have reported longer TTE and subsequently longer T@VO<sub>2max</sub> values for continuous exercises performed at intensities slightly less than iVO<sub>2max</sub> (92 – 95% iVO<sub>2max</sub>) than those performed at iVO<sub>2max</sub>.<sup>7 39</sup> T@VO<sub>2max</sub> appears to be influenced by TTAVO<sub>2max</sub> and TTE.<sup>19</sup> Hence, T@VO<sub>2max</sub> can be increased by reducing the TTAVO<sub>2max</sub> or increasing the TTE. The present results indicate that increasing the TTE may be more important for extending T@VO<sub>2max</sub>, since the faster TTAVO<sub>2max</sub> during CONT<sub>1</sub> did not result in a longer mean T@VO<sub>2max</sub>.

Comparison of T@VO<sub>2max</sub> between studies is limited due to differences in the method used to calculate T@VO<sub>2max</sub>. Several studies calculated T@VO<sub>2max</sub> by subtracting TTAVO<sub>2max</sub> from TTE.<sup>7 21 39</sup> This assumes that all VO<sub>2</sub> values from the time VO<sub>2max</sub> is elicited until exercise is terminated are  $\geq$  VO<sub>2max</sub>. It is believed that this method overestimates T@VO<sub>2max</sub> because as can be seen in figures 1 and 2, during exercise at iVO<sub>2max</sub>, individual VO<sub>2</sub> values can fall below

VO<sub>2max</sub> when VO<sub>2max</sub> has already been attained. Some studies have accepted data as being at VO<sub>2max</sub> when  $\geq 95\%$  VO<sub>2max</sub><sup>16 44</sup>. It has been established that T@VO<sub>2max</sub> is significantly longer when calculated using VO<sub>2</sub> data points that are  $\geq 95\%$  VO<sub>2max</sub> in comparison to VO<sub>2</sub> data points that are  $\geq$  VO<sub>2max</sub>.<sup>23</sup> Therefore, it may be more appropriate to compare T@VO<sub>2max</sub> recorded in these studies with T@95% VO<sub>2max</sub> values reported in the present study. Only one study by Hill *et al*<sup>39</sup> reported longer T@95% VO<sub>2max</sub> values ( $388 \pm 161$ ) than the present study, achieved during continuous exercise at 92% iVO<sub>2max</sub>.<sup>39</sup> The longer T@95% VO<sub>2max</sub> values resulted from a longer TTE ( $621 \pm 184$ ) than reported in the present study, since TTA95% VO<sub>2max</sub> was slower ( $233 \pm 92$ ) owing to the lower exercise intensity that was utilised. The results of Hill *et al*<sup>39</sup> and those of the present study further highlight the importance of prolonging TTE to maximise T@VO<sub>2max</sub> and T@95% VO<sub>2max</sub> during continuous exercise.

## Conclusions

In conclusion, the time spent exercising at a high level of VO<sub>2</sub> was increased when iVO<sub>2max</sub>(3 min) was utilised as the exercise intensity. TTE was longer during exercise at iVO<sub>2max</sub>(3 min) and this probably accounted for the longer T@VO<sub>2max</sub> and T@95% VO<sub>2max</sub> values derived during CONT<sub>3</sub>. Hence, if the goal is to maintain VO<sub>2max</sub>, iVO<sub>2max</sub>(3 min) is a more appropriate exercise intensity than iVO<sub>2max</sub>(1 min).

## REFERENCES

1. Daniels J, Scardina N, Hayes J, Foley, P. Elite and subelite female middle-and long distance runners. In: Landers DM, editor. Sport and elite performers: Proceedings of the 1984 Olympic Scientific Congress, July 19-23. Oregon: Champaign IL: Human Kinetics; 1984. p. 57-72.
2. Hill DW, Rowell AL. Running velocity at VO<sub>2max</sub>. Med Sci Sports Exerc 1996; 28:114-9.
3. Laursen PB, Jenkins DG. The scientific basis for high intensity interval training. Sports Med 2002; 32:53-73.

4. Wenger HA, Bell GJ. The interactions of intensity, frequency and duration of exercise training in altering cardiorespiratory fitness. *Sports Med* 1986; 3:346-356.
5. Berthon P, Fellmann N. General review of maximal aerobic velocity measurement at laboratory: proposition of a new simplified protocol for maximal aerobic velocity assessment. *J Sports Med Phys Fitness* 2002; 42:257-266.
6. Billat V, Renoux JC, Pinoteau J, Petit B, Koralsztein JP. Reproducibility of running time to exhaustion at  $VO_{2max}$  in subelite runners. *Med Sci Sports Exerc* 1994; 26:254-7.
7. Hill DW, Rowell AL. Responses to exercise at the velocity associated with  $VO_{2max}$ . *Med Sci Sports Exerc* 1997; 29:113-136.
8. Midgley AW, McNaughton LR. Time at or near  $VO_{2max}$  during continuous and intermittent running: a review with special reference to considerations for the optimisation of training protocols to elicit the longest time at or near  $VO_{2max}$ . *J Sports Med Phys Fitness* 2006; 46:1-14.
9. Millet GP, Candau R, Fattori P, Bignet, F.  $VO_2$  responses to different intermittent runs at velocity associated with  $VO_{2max}$ . *Can J Appl Physiol* 2003; 28:410-423.
10. Billat VL, Blondel N, Berthoin S. Determination of the velocity associated with the longest time to exhaustion at maximal oxygen uptake. *Eur J Appl Physio* 1999; 80:159-161.
11. Billat VL, Slawinski J, Bocquet V, Chassaing P, Demarle A, Koralsztein JP. Very short (15 s – 15 s) interval training around the critical velocity allows middle aged runners to maintain  $VO_{2max}$  for 14 minutes. *Int J Sports Med* 2001; 22:201–8.
12. Wakefield BR, Glaister M. Influence of work-interval intensity and duration on time spent at a high percentage of  $VO_{2max}$  during intermittent supramaximal exercise. *J Strength Cond Res* 2009; 23:2548 – 2554.
13. Zafeiridis A, Sarivasiliou H, Dipla K, Vrabas IS. The effects of heavy continuous versus long and short intermittent aerobic exercise protocols on oxygen consumption, heart rate, and lactate responses in adolescents. *Eur J Appl Physiol* 2010; 110:17–26.
14. Billat V, Faina M, Sardella F, Marini C, Fanton F, Lupo S, et al. A comparison of time to exhaustion at  $VO_{2max}$  in elite cyclists, kayak paddlers, swimmers and runner. *Ergonomics* 1996; 39:267–277.
15. Billat VL, Hill DW, Pintoneau J, Petit B, Koralsztein JP. Effect of protocol on determination of velocity at  $VO_{2max}$  and on its time to exhaustion. *Arch Physiol Biochem* 1996;104:313-321.
16. Dupont G, Blondel N, Lensele G, Berthoin S. Critical velocity and time spent at a high level of  $VO_2$  for short intermittent runs at supramaximal velocities. *Can J Appl Physiol* 2002;27:103-115.
17. Faina M, Billat V, Squadrone R, De Angelis M, Koralsztein JP, Dal Monte A. Anaerobic contribution to the time to exhaustion at the minimal exercise intensity at which maximal

- oxygen uptake occurs in elite cyclists, kayakists and swimmers. *Eur J Appl Physiol* 1997; 76:13–20.
18. Midgley AW, McNaughton LR, Wilkinson M. Criteria and other methodological considerations in the evaluation of time at  $VO_{2max}$ . *J Sports Med Phys Fitness*, 46, 183-188.
  19. Midgley AW, McNaughton LR, Wilkinson M. The relationship between the lactate turnpoint and the time at  $VO_{2max}$  during a constant velocity run to exhaustion. *Int J Sports Med* 2006;27:278-282.
  20. O'Brien BJ, Wibskov J, Knez WL, Paton CD, Harvey JT. The effects of interval – exercise duration and intensity on oxygen consumption during treadmill running. *J Sci Med Sport* 2008;11:287–290.
  21. Billat VL, Morton RH, Blondel N, Berthoin S, Bocquest V, Koralsztejn JP, et al. Oxygen kinetics and modelling of time to exhaustion whilst running at various velocities at maximal oxygen uptake. *Eur J Appl Physiol* 2000;82:178–187.
  22. Thevenet D, Leclair E, Tardieu-Berger M, Berthoin S, Regueme S, Prioux J. Influence of recovery intensity on time spent at maximal oxygen uptake during an intermittent session in young endurance trained athletes. *J Sports Sci* 2008;26:1313–1321.
  23. Midgley AW, McNaughton LR, Carroll S. Time at  $VO_{2max}$  during intermittent treadmill running: Test protocol dependent or methodological artefact? *Int J Sports Med* 2006;28:934-939.
  24. Robergs RA, Chwalbinska-Montea J, Mitchell JB, Pascoe DD, Houmard J, Costill DL. Blood lactate threshold differences between arterialised and venous blood. *Int J Sports Med* 1990;11:446–451.
  25. Thoden JS. (1991). Testing aerobic power. In: MacDougall JD, Wenger HA, Green HJ, editors. *Physiological testing of the high performance athlete*. Champaign IL: Human Kinetics; 1991. p. 107-175.
  26. Weltman A, Snead D, Stein P, Seip R., Schurrer R., Rutt R, et al. Reliability and validity of a continuous incremental treadmill protocol for the determination of lactate threshold, fixed blood lactate concentrations and  $VO_{2max}$ . *Int J Sports Med* 1990;11:26-32.
  27. Meyer T, Georg T, Becker C, Kindermann W. Reliability of gas exchange measurement from two different spiroergometry systems. *Int J Sports Med* 2001;22:593–7.
  28. Davison RRC, Coleman D, Balmer J, Nunn M, Theakston S, Burrows M, et al. Assessment of blood lactate: practical evaluation of the Biosen 5030 lactate analyzer. *Med Sci Sports Exerc* 2000;32:243–7.
  29. Heck H, Mader A, Hess G, Mucke, S, Muller R, Hollmann W. Justification of the 4-mmol/l lactate threshold. *Int J Sports Med* 1985;6:117–130.

30. Newell J, Higgins DN, Madden N, Cruikshank J, Einbeck J, McMillan K, et al.. Software for calculating blood lactate endurance markers. *J Sports Sci* 2007;25:1403-9.
31. James DVB, Sandals LE, Wood DM, Jones AM. Pulmonary gas exchange. In: Winter EM, Jones AM, Davison RRC, Bromley PD, Mercer TH, editors. *Sport and exercise physiology testing guidelines volume I: sport testing: The British Association of Sport and Exercise Sciences (BASES) guidelines*. London: Routledge; 2007. p. 101-111
32. Shapiro SS, Wilk MB. An analysis of variance test for normality (complete samples). *Biometrika* 1965;52:591–611.
33. Billat V, Renoux JC, Pinoteau J, Petit B, Koralsztein JP. Times to exhaustion at 90, 100 and 105% of velocity at  $VO_{2max}$  (maximal aerobic speed) and critical speed in elite longdistance runners. *Arch Phys Biochem* 1995;103:129–135.
34. Westerblad H, Allen DG. Recent advances in the understanding of skeletal muscle fatigue. *Curr Opin Rheumatol* 2002;14:648–652.
35. Westerblad H, Allen DG, Lannergren J. Muscle fatigue: lactic acid or inorganic phosphate the major cause? *News Physiol Sci* 2002;17:17–21.
36. Laursen PB, Shing CM, Jenkins DG. Reproducibility of the cycling time to exhaustion at  $VO_{2peak}$  in highly trained cyclists. *Can J Appl Physiol* 2003;28:605–615.
37. Caputo F, Denadai BS. Exercise mode affects the time to achieve  $VO_{2max}$  without influencing maximal exercise time at the exercise intensity associated with in triathletes. *Int J Sports Med* 2006;27:798–803.
38. Costa VP, De Matos DG, Pertence LC, Martins JAN, De Lima JRP. Reproducibility of cycling time to exhaustion at  $VO_{2max}$  in competitive cyclists. *J Exerc Physiol Online* 2011; 14:28–34.
39. Hill DW, Williams CS, Burt SE. Responses to exercise at 92% and 100% of the velocity associated with  $VO_{2max}$ . *Int J Sports Med* 1996;18:325-329.
40. Rossiter HB, Ward SA, Kowalchuk JM, Howe FA, Griffiths JR, Whipp BJ. Dynamic asymmetry of phosphocreatine concentration and  $O_2$  uptake between the on- and off-transients of moderate- and high-intensity exercise in humans. *J Physiol* 2002;15:991–1002.
41. Whipp BJ, Mahler M. Dynamics of gas exchange during exercise. In: West JB, editor. *Pulmonary gas exchange vol. II*. New York: Academic Press; 1980. p. 33–96.
42. Balaban RS. The role of  $Ca^{2+}$  signalling in the coordination of mitochondrial ATP production with cardiac work. *Biochim Biophys Acta* 2009;1787:1334-1341.
43. Chance B, Williams GR. Respiratory enzymes in oxidative phosphorylation. I. Kinetics of oxygen utilization. *J Biol Chem* 1955;217:383 - 393.

44. Billat VL, Bocquet V, Slawinski J, Laffite L, Demarle A, Chassaing P, et al. Effect of a prior intermittent run at  $v\dot{V}O_{2max}$  on oxygen kinetics during an all out severe run in humans. J Sports Med Phys Fitness 2000;40:185–194.

**Table 1.** Responses to a maximal incremental cycle test for the determination of  $i\dot{V}O_{2max}$  and  $\dot{V}O_{2max}$ . Values are means  $\pm$  standard deviations.

	$i\dot{V}O_{2max}$ (W)	$\dot{V}O_{2max}$ (L.min <sup>-1</sup> )	$\dot{V}O_{2max}$ (ml.kg <sup>-1</sup> .min <sup>-1</sup> )	$\dot{V}O_{2max}(DAY)$ (L.min <sup>-1</sup> )	$\dot{V}O_{2max}(DAY)$ (ml.kg <sup>-1</sup> .min <sup>-1</sup> )
$i\dot{V}O_{2max}(1 \text{ min})$	337 $\pm$ 36*	3.59 $\pm$	49.06 $\pm$	3.54 $\pm$	48.35 $\pm$
$i\dot{V}O_{2max}(3 \text{ min})$	292 $\pm$ 43	3.53 $\pm$	48.20 $\pm$	3.46 $\pm$	47.20 $\pm$


$\dot{V}O_{2max}$  = maximal oxygen uptake;  $i\dot{V}O_{2max}(1 \text{ min})$  = minimal exercise intensity to elicit  $\dot{V}O_{2max}$  during an incremental test with 1 min stages;  $i\dot{V}O_{2max}(3 \text{ min})$  = minimal exercise intensity to elicit  $\dot{V}O_{2max}$  during an incremental test with 3 min stages;  $\dot{V}O_{2max}(DAY)$  =  $\dot{V}O_{2max}$  of the day.  $\dot{V}O_{2max}$  determined during  $i\dot{V}O_{2max}(1 \text{ min})$  and  $i\dot{V}O_{2max}(3 \text{ min})$  tests.  $\dot{V}O_{2max}(DAY)$  determined during the  $i\dot{V}O_{2max}(1 \text{ min}) + 20 \text{ W}$  and  $i\dot{V}O_{2max}(3 \text{ min}) + 20 \text{ W}$  tests. \*Indicates significant differences.

**Table 2.** Physiological responses to a continuous bout of cycling to exhaustion at  $i\dot{V}O_{2max}(1 \text{ min})$  (CONT<sub>1</sub>) and a second continuous bout of cycling to exhaustion at  $i\dot{V}O_{2max}(3 \text{ min})$  (CONT<sub>3</sub>). Values are means  $\pm$  standard deviations.


	TTE (s)	T@ $\dot{V}O_{2max}$ (s)	T@95% $\dot{V}O_{2max}$ (s)	TTA $\dot{V}O_{2max}$ (s)	TTA95% $\dot{V}O_{2max}$ (s)	BLa (mMol.L <sup>-1</sup> )	HR <sub>max</sub> (beats.min <sup>-1</sup> )
CONT <sub>1</sub>	214 $\pm$ 65	11 $\pm$ 20	57 $\pm$ 51	185 $\pm$ 54	97 $\pm$ 34	10.8 $\pm$ 1.5	188 $\pm$ 10
CONT <sub>3</sub>	529 $\pm$ 140*	146 $\pm$ 158*	326 $\pm$ 211*	407 $\pm$ 118*	173 $\pm$ 76*	9.0 $\pm$ 1.2*	189 $\pm$ 10
Mean	315	134	269	222	76	1.8	1
diff	[209-424]	[8-260]	[110-428]	[144-299]	[10-142]	[1.1-2.5]	[0-1]
[95% CL]							

CONT<sub>1</sub> = continuous cycling to exhaustion at  $i\dot{V}O_{2max}(1 \text{ min})$ ; CONT<sub>3</sub> = continuous cycling to exhaustion at  $i\dot{V}O_{2max}(3 \text{ min})$ ; TTE = time to exhaustion; T@ $\dot{V}O_{2max}$  = time at  $\dot{V}O_{2max}$ ; T@95% $\dot{V}O_{2max}$  = time at 95%  $\dot{V}O_{2max}$ ; TTA $\dot{V}O_{2max}$  = time to achieve  $\dot{V}O_{2max}$ ; TTA95% $\dot{V}O_{2max}$  = time to achieve 95%  $\dot{V}O_{2max}$ ; BLa = blood lactate concentration; HR<sub>max</sub> = maximum heart rate; Mean diff = mean difference between CONT<sub>1</sub> and CONT<sub>3</sub>; 95% CL = 95% confidence limits; \*Indicates significant differences between mean values.





**Figure 1.** Example of the oxygen uptake response observed in a single participant during a continuous bout of cycling to exhaustion at the lowest exercise intensity associated with  $VO_{2max}$  during an incremental test with one minute stages (20 W increments). Individual data points are 10 s average values.  $TTAVO_{2max}$  = time to achieve  $VO_{2max}$ ;  $T@VO_{2max}$  = time at  $VO_{2max}$ .



**Figure 2.** Example of the oxygen uptake response observed in a single participant during a continuous bout of cycling to exhaustion at the lowest exercise intensity associated with  $VO_{2max}$  measured during an incremental test with three minute stages (20 W increments). Individual data points are 10 s average values.  $TTAVO_{2max}$  = time to achieve  $VO_{2max}$ ;  $T@VO_{2max}$  = time at  $VO_{2max}$ .