

**The influence of compression garments on recovery following Marathon running**

Jessica Hill<sup>1</sup>, Glyn Howatson<sup>2,4</sup>, Ken van Someren<sup>3</sup>, Ian Walshe<sup>2</sup> and Charles Pedlar<sup>1</sup>

<sup>1</sup>*School of Sport, Health, and Applied Science, St. Mary's University College, Twickenham, UK.* <sup>2</sup>*School Life of Sciences, Northumbria University, Newcastle Upon Tyne, UK.* <sup>3</sup>*English Institute of Sport, Marlow, UK.* <sup>4</sup>*Centre for Aquatic Research, University of Johannesburg, South Africa.*

*Corresponding author: Jessica Hill, School of Sport, Health, and Applied Science, St. Mary's University College, Twickenham, TW1 4SX UK. Tel: +00 44 0 208 240 4000, Fax: +00 44 0 208 240 4212, Email [jessica.hill@smuc.ac.uk](mailto:jessica.hill@smuc.ac.uk).*

**Running Head**

Compression garments and Marathon recovery

ACCEPTED

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27

**ABSTRACT**

Strenuous physical activity can result in exercise induced muscle damage. The purpose of this study was to investigate the efficacy of a lower limb compression garment in accelerating recovery from a marathon run. Twenty four subjects (female n= 7, male n= 17) completed a marathon run before being assigned to a treatment group or a sham treatment group. The treatment group wore lower limb compression tights for 72 h following the marathon run, the sham treatment group received a single treatment of 15 min of sham ultrasound following the marathon run. Perceived muscle soreness, maximal voluntary isometric contraction (MVIC) and serum markers of Creatine Kinase (CK) and C-reactive protein (C-RP) were assessed before, immediately after and 24, 48 and 72 h post marathon. Perceived muscle soreness was significantly lower ( $p < 0.05$ ) in the compression group at 24 h post marathon when compared to the sham group. There were no significant group effects for MVIC, CK and C-RP ( $p > 0.05$ ). The use of a lower limb compression garment improved subjective perceptions of recovery, however there was no significant improvement in muscular strength, nor was there a significant attenuation in markers of exercise induced muscle damage and inflammation.

**Key Words:** Exercise, Muscle damage, Endurance, External pressure

1 **INTRODUCTION**

2

3 Exercise induced muscle damage (EIMD) often occurs following strenuous physical activity, particularly when  
4 the exercise has an eccentric component or is of a prolonged duration, both of these characteristics are typical of  
5 long distance running (7, 29). This phenomenon is characterised by a number of symptoms including pain,  
6 inflammation and reduced capacity of the affected muscle to produce force (10, 12). The negative symptoms  
7 associated with muscle damage are likely to affect the functional capability of the individual, resulting in a  
8 reduction in subsequent exercise performance (13). Strategies that may reduce symptoms of EIMD and enhance  
9 the recovery process are desirable, to enable the athlete to tolerate a higher training volume, frequency or  
10 intensity (8).

11

12 Prolonged endurance running is reported to cause damage to the muscle fiber and associated structures, this  
13 damage arises from several contributing processes (5). The initial damage is considered to be mechanical in  
14 nature resulting from repetitive loading on the muscle fibers (6). Following the initial mechanical damage, a  
15 secondary inflammatory process arises that is accompanied by oxidative stress. The experience of oxidative  
16 stress can further exacerbate the initial mechanical stress in a so called vicious cycle (4, 22). Elevations in  
17 markers of muscle damage and inflammation have been observed following a marathon run (24), confirming that  
18 the event causes significant ultrastructural disruption. It may therefore be possible to apply an intervention that  
19 can effectively manage the processes associated with secondary damage, either by reducing the magnitude of  
20 damage or by accelerating the recovery process.

21

22 Knowledge regarding the potential mechanisms underpinning the efficacy of compression is limited, however it  
23 is suggested that compression garments may be effective in reducing the swelling and inflammatory processes  
24 associated with muscle damage (21). It is suggested that the garment works by creating an external pressure  
25 gradient that reduces the space available for swelling to occur, thereby reducing the secondary inflammatory  
26 response (13). It has also been suggested that the use of compression can improve venous return, reduce venous  
27 pooling and enhance the removal of metabolites, this is perhaps due to an enhanced muscle pump function (3,  
28 26). Previous research has observed reduced perceptions of muscle soreness; (3, 13), reduced concentrations of  
29 serum creatine kinase (CK) (3, 13) and a reduced inflammatory response (17, 28) with the use of compression

1 garments. In contrast, research has also demonstrated that compression garments have no effect on muscle  
2 soreness (2, 17), CK concentration (2, 13), lactate dehydrogenase concentration (13), and thigh girth (13).  
3 Although the physiological and biochemical effects of wearing compression garments remain poorly understood,  
4 the positive claims associated with the garments may have led to an increase in the use of compression garments  
5 in sport.

6 Differences in experimental design including exercise modality, duration of the application of compression and  
7 the participant's training status may explain some of the inconsistencies within the literature. Furthermore, the  
8 majority of studies have focused on power sports (14, 15, 27). Participation in endurance events such as  
9 marathon running is increasing in popularity, and the use of compression garments as training aids is also  
10 increasing (14). Thus there is a need to focus on the efficacy of compression garments on recovery in endurance  
11 based sports. To the knowledge of the authors, the use of compression following prolonged endurance running  
12 has not been examined. Accordingly the purpose of this study was to investigate the effects of wearing a  
13 commercially available, lower limb, compression garment on the recovery of strength, soreness and indices of  
14 muscle damage following a marathon run.

15

## 16 ***MATERIALS AND METHODS***

### 17 ***Subjects***

18 Following approval from the University's research ethics committee, in accordance with the Declaration of  
19 Helsinki, 24 recreational marathon runners (male; n=17; female: n=7), volunteered to participate in this  
20 investigation. All subjects completed a health screening questionnaire and gave written informed consent.  
21 Participant characteristics, marathon time and marathon history are presented in table 1. Subjects were asked to  
22 refrain from heavy exercise in the 48 h preceding the marathon. Subjects were also asked to refrain from using  
23 any recovery strategy, such as water immersion, NSAIDs, massage and also to refrain from consuming alcohol  
24 for the duration of the study.

25

### 26 ***Experimental Design***

27 Subjects were required to complete a 26.2 mile self paced run in an outside environment. Subjects could select  
28 when to complete the marathon on one of three dates, the course remained the same on each date. The mean

1 environmental conditions for the three days were a barometric pressure of  $762 \pm 12$  mmHg (750-771 mmHg), an  
2 air temperature of  $9 \pm 2^\circ\text{C}$  (6-10° C), a wind speed of  $16 \pm 4$  km·h<sup>-1</sup> (12-20 km·h<sup>-1</sup>) and a relative humidity of 73  
3  $\pm 6\%$  (66-77%). At least 7 days before the marathon run subjects reported to the laboratory to complete a  
4 submaximal lactate profile and a maximal exercise test on a treadmill (H/P Cosmos Pulsar 4.0, H/P Cosmos  
5 Sports and Medical, Nussdorf-Traunsdein, Germany), in order to characterise the physiology of the group and  
6 establish baseline data. The submaximal lactate profile consisted of a series of stages each lasting 3 min with 30  
7 s rest between each stage to enable the collection of a blood lactate sample, the sample was immediately  
8 analysed using an automated analyser (Biosen C-Line, EKF Diagnostic, Ebendorfer Chaussee 3, Germany). At  
9 the start of each new stage the treadmill speed was increased by 1 km·h<sup>-1</sup>, heart rate (Polar A1, Polar Electro OY,  
10 Kempele, Finland) was recorded throughout the test and a 1 min average was taken during the last minute of  
11 each stage. Ratings of perceived exertion (RPE) were assessed using the Borg scale. Subjects were asked to  
12 indicate their perception of exertion on a scale ranging from 6 (very, very light) to 20 (very, very hard) during  
13 the last 30 s of each stage. Expired air was analysed continuously using an online gas analyser (Jaeger Oxycon  
14 Pro, Jaeger Ltd, Hoechberg, Germany). The submaximal test was terminated when blood lactate concentration  
15 exceeded 4 mmol.L<sup>-1</sup>. The participant was given 10 min rest before commencing the maximal exercise test. The  
16 test began 2 km·h<sup>-1</sup> slower than the finishing speed of the submaximal test. Throughout the test the treadmill  
17 speed remained constant and the gradient was increased by 1% every minute. The test was terminated when the  
18 participant reached volitional exhaustion. Expired air was analysed continuously, maximal oxygen uptake  
19 ( $\dot{V}O_{2\text{max}}$ ) was determined as the highest 30 s average. Verbal encouragement was given throughout. Subjects  
20 were matched based upon  $\dot{V}O_{2\text{max}}$  and then were randomly assigned to a compression garment group  
21 (compression) or a sham ultrasound group (sham). Dependent measures included perceived muscle soreness,  
22 maximum voluntary isometric contraction (MVIC) of the knee extensors, creatine kinase (CK) and C-reactive  
23 protein (C-RP) and were assessed immediately before the marathon, within 1 h post marathon and at 24, 48 and  
24 72 h post marathon.

25

## 26 ***Treatment Groups***

27 The sham group received 15 min (5 min quadriceps, 5 min, hamstrings and 5 min gastrocnemius) of sham  
28 ultrasound (Combined therapy ultrasound/inferential, Shrewsbury Medical, Shropshire, UK) within 1 h of the  
29 completion of the marathon run. A water-soluble hypoallergenic ultrasound gel (Aquasonic 100 ultrasound

1 transmission gel, Parker Laboratories, Fairfield, USA) was applied to the leg and the ultrasound head was used  
2 to spread the gel over the skin in circular patterns. The ultrasound unit was turned off throughout the duration  
3 and the unit was obscured from view. The compression garment group were given a pair of compression tights  
4 (2XU, MA1551b men's compression tights and WA1552b women's compression tights, Melbourne, Australia),  
5 fitted according to the manufacturer's instructions. Subjects were instructed to shower and put on the garments  
6 within 1 h of the completion of the marathon run. The garments were worn for 72 h following the run, subjects  
7 were asked to wear them continuously, only removing them to shower. The degree of pressure exerted on the  
8 leg was measured using a body pressure-measuring device (Kikuhime, TT Medi Trade, Søleddet, Denmark).  
9 Pressure was measured at the medial aspect of the calf at the site of maximal girth and also at the front thigh at  
10 the midpoint between the inguinal crease and the superior aspect of the patella. Pressure measurements at each  
11 site were taken with the subject standing in the anatomical position and also during a contraction of each muscle  
12 group. Measurements were repeated 3 times with the mean value recorded.

13

#### 14 *Dependent Variables*

15 Muscle soreness was obtained using a 200 mm visual analogue scale with 'no pain' at 0 mm and 'unbearable  
16 pain' at 200 mm (Howatson et al. 2010). Subjects stood with their feet shoulder width apart with hands on hips  
17 and were asked to perform a squat to 90° and return to standing and mark their subjective feeling of pain on the  
18 scale. MVIC was assessed using a strain gauge (MIE Medical Research Ltd., Leeds, UK). Subjects were seated  
19 on a platform, with their hip and knee joints flexed at 90°. The strain gauge was attached 2 cm above the  
20 malleoli of the left ankle. Subjects were asked to maximally extend the knee against the device for 3 s. Each  
21 participant was given 3 maximal attempts each separated by 1 min the highest of which was recorded as MVIC.  
22 Verbal encouragement was given (25).

23

24 Serum CK and C-RP were measured at each time point. Approximately 8.5 mL of blood was collected from the  
25 antecubital fossa into serum separation vacutainers (BD equipment, vacutainer systems, Plymouth, UK).  
26 Following collection the sample was immediately placed in a refrigerated centrifuge (Mikro 220R D-78532,  
27 tuttlingen, Germany) and spun at 3500 rpm, a relative centrifugal force of 3000g, for 20 minutes at 4°C to enable  
28 the separation of serum. The serum was immediately frozen at -80°C for later analysis.

29

1 Creatine kinase was analysed in duplicate using an automated clinical chemistry analyser (Modular P, Roche  
2 Diagnostics, West Sussex, UK). C-RP was analysed in duplicate from serum samples using a high sensitivity C-  
3 reactive protein ELISA kit (Kalon Biological, Guildford, UK). The absorbance of each sample was read at  
4 405nm (Biochrom Ltd, Cambridge, UK).

5

## 6 *Statistical Analysis*

7 All data analyses were carried out using SPSS for Windows v 18 and values are reported as mean  $\pm$  SD. An  
8 independent samples t-test was used to identify any differences in group characteristics at baseline. In order to  
9 account for inter-individual variation MVIC was expressed as a percentage change from baseline. All dependent  
10 variables were assessed using a treatment by time repeated measures analysis of variance (ANOVA). Mauchly's  
11 test of sphericity was used to check homogeneity of variance for all analysis. The Greenhouse-Geisser  
12 correction factor was used where violations of the assumption occurred. Where a significant effect was observed  
13 interaction effects were further examined using an LSD *post hoc* analysis. A significance level of  $p \leq 0.05$  was  
14 applied.

15

## 16 **RESULTS**

17 The mean pressure of the garment exerted on the front thigh was  $9.9 \pm 2$  mmHg in the anatomical position and  
18  $17.5 \pm 3.5$  mmHg during muscle contraction. The mean pressure exerted on the calf was  $19.3 \pm 2.6$  mmHg in the  
19 anatomical position and  $24.4 \pm 5$  mmHg during muscle contraction. There were no significant differences in  
20  $\dot{V}O_{2\max}$ , previous marathon history, average weekly training distance and marathon finish time between groups  
21 (Table 1). All dependent variables showed a significant time effect ( $p \leq 0.004$ ).

22

23 Muscle soreness exhibited a significant time by group effect ( $F_{(4,1)} = 30.8, p < 0.001$ ) and a significant group  
24 effect ( $F_{(1,22)} = 4.451, p = 0.046$ , with those in the compression group experiencing reduced muscle soreness  
25 (Figure 1). A *post hoc* LSD test indicated that this difference occurred at 24 h post exercise in the compression  
26 group. At this time point the sham group increased in mean soreness from 26.8 to 36.4 mm, while the  
27 compression group decreased in mean soreness from 31 to 13.9 mm.

28

1 Post marathon concentrations for CK were elevated at 24 and 48 h post exercise, with the highest values  
2 occurring at 24 h ( $692.1 \pm 625.6$  and  $1022.3 \pm 1439$  U/L in the compression and sham group respectively). C-  
3 RP concentrations were also elevated at 24 and 48 h post marathon, with the highest values occurring at 24 h  
4 post ( $7.4 \pm 4.6$  and  $10.3 \pm 9.0$   $\text{mg}^{-1}$  in the compression and sham group respectively), however there was no  
5 significant difference between groups for either marker ( $p \geq 0.05$ ; see figures 2 and 3).

6

7 Maximal voluntary isometric contraction (MVIC) was reduced immediately post marathon, returning to normal  
8 at 72h h post marathon. The largest measured decrease in MVIC occurred immediately after the marathon ( $82.8$   
9  $\pm 13.9$  and  $77 \pm 14.8$  percent of max in the compression and sham group respectively), however no significant  
10 group effect ( $F_{(1,22)} = 2.256$ ,  $p = 0.124$ ) or interaction ( $F_{(4,1)} = 0.688$ ,  $p = 0.543$ ) was observed (see figure 4).

11

## 12 **DISCUSSION**

13 To the authors knowledge this was the first investigation to examine the efficacy of compression garments in  
14 accelerating recovery from endurance running. It was hypothesised that the use of lower limb compression  
15 garment would enhance the recovery process following a marathon run. Those in the compression garment  
16 group experienced less muscle soreness 24 h post marathon when compared to the sham group. Despite this  
17 finding there were no significant differences between groups for MVIC, CK and C-RP, however there appears to  
18 be a trend for improved recovery of MVIC and an attenuation of CK and C-RP in the compression group.

19

20 The main finding of the present study was that the use of a lower limb compression garment significantly  
21 reduced muscle soreness 24 h after a marathon run (see Figure 1). This finding is consistent with other studies  
22 which have demonstrated that the use of compression garments results in reduced perceived muscle soreness  
23 following damaging exercise (3, 13). The experience of muscle soreness is proposed to arise from mechanical  
24 trauma to the tissue, giving rise to structural damage and an inflammatory response (11). Therefore a reduction  
25 in soreness may be related to reduced structural damage to the tissue and/or a reduced inflammatory response  
26 (3). It is suggested that the application of compression may be effective in reducing this inflammatory process,  
27 this may be because compression garments create an external pressure gradient reducing the space available for  
28 swelling, haemorrhage and haematoma (13, 21). The chain of events that occur following initial mechanical  
29 damage to the muscle fibers can result in an increased osmotic pressure within the tissue, which causes an efflux



1 of fluid from the capillaries into the interstitial spaces (21). The application of compression may lessen the  
2 change in osmotic pressure and consequently reduce inflammation, the circulation of inflammatory markers and  
3 the sensation of pain (21). Although this study controlled for a placebo effect by using sham ultrasound, it is  
4 possible that the finding of reduced perceived muscle soreness may be linked to the participant's belief that  
5 compression garments have a positive response on recovery.

6

7 C-Reactive protein is a marker of inflammation that has been proposed to reflect the inflammatory load on the  
8 body (30). Concentrations of C-RP were elevated following the marathon run, peaking at 24 h post marathon  
9 (see Figure 2), this is consistent with previous research (24). There was no significant group effect observed on  
10 concentrations of C-RP. Few studies investigating the use of compression garments have measured C-RP, those  
11 who have measured this marker have also found no significant effect of compression on concentrations (16).  
12 However, several have looked at mid thigh girth as an indirect measure of inflammation, one such study  
13 observed no time or group effect with mid thigh girth following the application of compression (13). This data  
14 differs from the findings of an earlier study that observed an increased thigh circumference in a contrast bathing  
15 group and a control group but no increase in thigh circumference in a compression garment group, following a  
16 bout of damaging exercise (17). Given that there were no significant reductions in the inflammatory marker  
17 CRP, the reduction of soreness is unlikely to be linked to a reduction in the inflammatory response.

18

19 Similarly to C-RP, this study observed no group effect for CK. Creatine kinase (CK), an enzyme that leaks from  
20 damaged muscle tissue into the blood stream resulting in an elevated serum concentration of CK (9), is  
21 frequently used as a marker of muscle damage. A peak increase in concentration is usually observed 24h post  
22 exercise with levels remaining elevated at 48 and 72 h post-exercise (9). The CK concentrations observed in this  
23 study follow the well established response to damaging exercise. The concentration of CK was elevated  
24 immediately after the marathon run, peaking at 24 h post exercise. The concentration of CK continued to  
25 remain elevated at 48 and 72 h post-exercise (see Figure 3). This indicates that the marathon run induced a  
26 significant amount of tissue damage. While it appears that there was a trend for higher concentrations of CK in  
27 the sham group this difference was not statistically significant. This apparent trend is consistent with the findings  
28 of previous studies that have found reduced concentrations of CK in those wearing compression garments  
29 compared to those in a control group (13, 14, 18).

1 Training status is a key factor that may explain why the current study has not observed a reduction in CK or C-  
2 RP when compared to previous research. Well trained individuals tend to exhibit lower concentrations of CK  
3 following a bout of damaging exercise when compared to their less trained counterparts (9). This is a reported  
4 benefit of the repeated bout effect which suggests that a bout of damaging exercise offers protection against  
5 future bouts of damaging exercise whereby the individuals experience less severe symptoms of EIMD (23). The  
6 average number of previous marathons completed by the subjects taking part in this study was  $16 \pm 24$   
7 suggesting that they were experienced marathon runners, it is therefore possible that the prior marathon and  
8 training history of the subjects in this study offered protection in the form of an adaptive response.

9  
10 Previous studies that have indicated a lowered CK concentration following the application of compression have  
11 used club level (14) or university level (13) games players who may not be familiar with high training volumes.  
12 Howatson et al. (24) observed much higher values of CK (2814 compared to 1022 U/L in the placebo groups)  
13 and CRP (27 compared to 10 mg/L in the placebo groups) following a marathon race compared to this present  
14 study. Higher values of these markers would indicate a greater degree of muscle damage occurred, this could be  
15 related to previous marathon history as the average number of competed marathons for the subjects taking part in  
16 Howatson et al. (24) was  $6 \pm 8$  where the average number of completed marathons for the subjects taking part in  
17 this study was  $16 \pm 24$ , indicating that the subjects in the present study were more experienced. It is possible that  
18 the potential of the garment to have a beneficial effect is attenuated with a lower magnitude of change in markers  
19 of muscle damage and inflammation. This is perhaps why, in the present study, there was a trend for reduced  
20 concentrations of CK and C-RP but no significant effect.

21  
22 The use of compression garments did not improve the recovery of muscle force production following a marathon  
23 run (see Figure 4). This is consistent with the findings of others (14, 16). The effects of compression on  
24 contractile performance is unclear, it is thought that any improvement or recovery in muscle strength may be  
25 related to contractile function. The decrease in MVIC observed immediately following the marathon run may be  
26 due to both localised muscle trauma and peripheral muscle fatigue, resulting from the accumulation of  $H^+$  ions  
27 which will inhibit excitation-contraction coupling (19). The results of this study indicate that, despite the use of  
28 compression, the recovery of voluntary muscle function was not improved. It has previously been suggested that  
29 equivocal responses in performance standards following the use of a recovery strategy may be due to training

1 status (17). Experienced athletes are more accustomed to the discomfort associated with high intensity exercise  
2 and are better able to reproduce performances (17).

3

4 One current argument surrounding the use of commercially available compression garments is whether they can  
5 exert enough pressure to be of any benefit (13). In contrast to this research has also suggested that garments  
6 exerting high levels of compression (23-32 mmHg) are not as effective at maintaining leg power as garments  
7 exerting low (12-15 mmHg) or medium (18-21 mmHg) pressures (1). The pressures observed within this study  
8 ranged between 9.9-24.4 mmHg. The fitting of these garments is often based upon the manufacturer's  
9 guidelines, however due to differences in body size and tissue structure the range of sizes available may not be  
10 sufficient. Whilst we measured the degree of compression exerted in this study and our compression level was  
11 similar to that achieved in other studies (3, 20), more research is needed to ascertain the optimal degree of  
12 compression in the recovery from symptoms of EIMD as well as how the degree of compression varies between  
13 individuals wearing the same size garments.

14

15 In conclusion, it would appear that the use of a lower limb compression garment, following a marathon run,  
16 results in significantly lower perceived muscle soreness at 24 h post marathon. In addition the garment had no  
17 statistically significant effect on improved recovery of muscle performance or post-exercise removal of serum  
18 markers of muscle damage and inflammation. However, despite the lack of a statistically significant effect in  
19 these markers a trend for improvement was apparent in the compression group.

20

### 21 ***Practical Application***

22 A variety of modalities are frequently used to accelerate recovery from strenuous exercise. A number of  
23 investigations have indicated that the use of compression garments may help recovery following damaging  
24 exercise (3, 13, 21). The observation of reduced muscle soreness in this study may have implications for a  
25 number of populations, wishing to embark on exercise regimes, who are discouraged due to the experience of  
26 soreness. As such compression garments may be of use as a recovery aid when worn during the 24 h post  
27 exercise to reduce the experience of perceived muscle soreness. While this study investigated the effects of

1 compression in experienced, but nonetheless recreational runners, further research should look to identify  
2 whether significantly greater effects are observed in less experienced runners

3

#### 4 ***Acknowledgements***

5 The authors would like to thank the subjects for their commitment throughout the completion of this project. We  
6 would also like to thank Rik Mellor, Hayley Legg, Hayley Morgan, Kylie Ilston, Nicola Brown, Natasha Mistry,  
7 Ammi King, Mel Bradley, Hilary Davies, John Pattison, Sunny Pottay, Steve Hiorns, Marcus Docker, Ross  
8 MacGregor and Richard Burden for their valuable help on the day of the marathon. Finally we would like to  
9 thank the St Mary's University College School Research Fund for supporting this project. This research  
10 investigation was carried out in compliance with the ethical laws of Great Britain and the authors have no  
11 conflicts of interest that are relevant to the content of this article. The results of the present study do not  
12 constitute endorsement of the product by the authors or the NSCA.

13

#### 14 **REFERENCES**

- 15 1. Ali, A, Creasy, RH, and Edge. The effect of graduated compression stockings on running  
16 performance. *J Strength Cond Res* 25: 1385-1392, 2011.
- 17 2. Ali, A, Creasy, RH, and Edge, JA. Physiological effects of wearing graduated compression  
18 stockings during running. *Eur J Appl Physiol* 109: 1017-1025, 2010.
- 19 3. Ali, A, Caine, MP, and Snow, BG. Graduated compression stockings: Physiological and perceptual  
20 responses during and after exercise. *J Sports Sci* 25: 413-419, 2007.
- 21 4. Aoi, W, Naito, Y, Takanami, Y, Kawai, Y, Sakuma, K, Ichikawa, H, Yoshida, N, and Yoshikawa, T.  
22 Oxidative stress and delayed-onset muscle damage after exercise. *Free Radic Biol Med* 37: 480-487,  
23 2004.
- 24 5. Appell, HJ, Soares, JMC, and Duarte. JAR. Exercise, muscle damage and fatigue. *Sports Med*  
25 13:108-115, 1992.
- 26 6. Armstrong, RB. Muscle damage and endurance events. *Sports Med* 3: 370-381, 1986.
- 27 7. Armstrong, RB. Initial events in exercise – induced muscular injury. *Med Sci Sports Exerc* 22: 429-435,  
28 1990.


- 1 8. Barnett, A. Using recovery modalities between training sessions in elite athletes. Does it help? *Sports*  
2 *Med* 36: 781-796, 2006.
- 3 9. Brancaccio, P, Maffulli, N, and Limongelli, FM. Creatine kinase monitoring in sport medicine. *Brit Med*  
4 *Bull* 81: 1-22, 2007.
- 5 10. Cleak, MJ, and Eston, RG. Muscle soreness, swelling, stiffness and strength loss after intense eccentric  
6 exercise. *Brit J Sport Med* 26: 267-271, 1992.
- 7 11. Connolly, DAJ, Sayers, SP, and McHugh MP. Treatment and prevention of delayed onset muscle  
8 soreness. *J Strength Cond Res* 17: 197-208, 2003.
- 9 12. Connolly, DAJ, Lauzon, C, Agnew, J, Dunn, M, and Reed, B. The effects of vitamin C supplementation  
10 on symptoms of delayed onset muscle soreness. *J Sport Med Phys Fit* 46: 462-467, 2006.
- 11 13. Davies, V, Thompson, KG, and Cooper, SM. The effects of compression garments on recovery. *J*  
12 *Strength Cond Res* 2: 1786-1794, 2009.
- 13 14. Doan, BK, Kwon, Y, Newton, RU, Shim, J, Popper, EM, Rogers, RA, Bolt, LR, Robertson, M, and  
14 Kraemer, WJ. Evaluation of a lower body compression garment. *J Sports Sci* 21: 602-610, 2003.
- 15 15. Duffield, R, and Portus, M. Comparison of three types of full body compression garments on throwing  
16 and repeat-sprint performance in cricket players. *Br J Sports Med* 41: 409-414, 2007.
- 17 16. Duffield, R, Cannon, J, and King M. The effects of compression garments on recovery of muscle  
18 performance following high-intensity sprint and plyometric exercise. *J Sci Med Sport* 13: 136-140,  
19 2010.
- 20 17. French, DN, Thompson, KG, Garland, SW, Barnes, CA, Portas, MD, Hood, PE, and Wilkes, G. The  
21 effects of contrast bathing and compression therapy of muscular performance. *Med Sci Sports Exerc* 40:  
22 1297-1306, 2008.
- 23 18. Gill, ND, Beaven, CM, and Cook, C. Effectiveness of post match recovery strategies in rugby players.  
24 *Br J Sports Med* 40: 260-263, 2006.
- 25 19. Green, HJ. Mechanisms of muscle fatigue in intense exercise. *J Sports Sci* 15: 247-256, 1997.
- 26 20. Kraemer, WJ, Volek, JS, Bush, JA, Gotshalk, LA, Wagner, PR, Gómez, AL, Zatsiorsky, VM, Duzrte,  
27 M, Ratamess, NA, Mazzetti, SA, and Selle, BJ. Influence of compression hosiery on physiological  
28 responses to standing fatigue in women. *Med Sci Sports Exerc* 32: 1849-1858, 2000.
- 29 21. Kraemer, WJ, French, DN, and Spiering, BA. Compression in the treatment of acute muscle injuries in  
30 sport. *Int Sportmed J* 5: 200-208, 2004.

- 1 22. Halliwell, B, and Chirico, S. Lipid peroxidation: its mechanism, measurement, and significance. *Am J*  
2 *Clin Nutr* 57: 715S-725S, 1993.
- 3 23. Howatson, G, and van Someren, K. The prevention and treatment of exercise induced muscle damage.  
4 *Sports Med* 38: 483-503, 2008.
- 5 24. Howatson, G, McHugh, MP, Hill, JA, Brouner, J, Jewell, AP, van Someren, KA, Shave, RE, and  
6 Howatson, SA. Influence of tart cherry juice on indices of recovery following marathon running. *Scan J*  
7 *Med Sci Spor* 20: 843-852, 2010.
- 8 25. Howatson, G, Hough, P, Pattison, J, Hill, JA, Blagrove, R, Glaister, M, and Thompson, KG. Trekking  
9 poles reduce exercise-induced muscle injury during mountain walking. *Med and Sci Sports Exerc* 43:  
10 140-145, 2011.
- 11 26. Ibegbuna, V, Delis, KT, Nicolaides, AN, and Aina, O. Effect of elastic compression stockings on  
12 venous hemodynamics during walking. *J Vasc Surg* 37: 420-425, 2003.
- 13 27. Montgomery, PG, Pyne, DB, Cox, AJ, Hopkins, WG, Dorman, JC, Cook, K, and Minahan, CL. The  
14 effect of recovery strategies on physical performance and cumulative fatigue in competitive basketball.  
15 *J Sports Sci* 26: 1135-1145, 2008.
- 16 28. Trenell, MI, Rooney, KB, Sue, CM, and Thompson, CH. Compression garments and recovery from  
17 eccentric exercise: A P-MRS study. *J Sports Sci Med* 5: 106-114, 2006.
- 18 29. Vickers, A.J. (2001). Time course of muscle soreness following different types of exercise. *BMC*  
19 *Musculoskel Dis* 2: 1-4, 2001.
- 20 30. Yeh, ETH. CRP as a mediator of disease. *Circulation* 109: 11-14, 2004.

21

**Table 1.** Descriptive characteristics of the participants in the compression and ultrasound groups. Values are mean  $\pm$  SD, N=12 per group. There were no statistically significant differences between groups for any variable.


<b>Group</b>	<b>Age (yrs)</b>	<b>Height (cm)</b>	<b>Mass (kg)</b>	<b>VO2max (ml·kg<sup>-1</sup>·min<sup>-1</sup>)</b>	<b>Finish time (h:min:s)</b>	<b>Previous Marathons</b>	<b>Weekly mileage</b>
<b>Compression</b> (n = 8 male, n = 4 female)	47.7 $\pm$ 10.8	177.8 $\pm$ 10.2	73.3 $\pm$ 14.1	53.8 $\pm$ 10.2	03:46:45 $\pm$ 00:22:30	15.6 $\pm$ 28.8	35.3 $\pm$ 10.5
<b>Sham ultrasound</b> (n = 9 male, n = 3 female)	41.1 $\pm$ 10.5	175.3 $\pm$ 7.0	71.6 $\pm$ 7.3	55.6 $\pm$ 8.4	03:39:27 $\pm$ 00:33:10	12.7 $\pm$ 12.1	40.9 $\pm$ 15.2




**Fig. 1** Perceived rating of muscle soreness for the compression and sham groups before and after the Marathon.

\* significantly different from compression group (grey lines denote individual participant results, black line denotes the group mean with SD values).






**Fig. 2** Serum CRP values for the compression and sham groups before and after the Marathon. No significant differences were observed between treatment groups (grey lines denote individual participant results, black line denotes the group mean with SD values).



**Fig. 3** Serum CK values for the compression and sham groups before and after the Marathon. No significant differences were observed between treatment groups (grey lines denote individual participant results, black line denotes the group mean with SD values).



**Fig. 4** Percentage change in MVIC for the compression and sham groups before and after the Marathon. No significant differences were observed between treatment groups (grey lines denote individual participant results, black line denotes the group mean with SD values).