

TITLE

Impact of Fixture Congestion on Indices of Performance & Recovery in Youth Soccer Players

AUTHOR

Hattersley, Chris; Wells, Carl; Blagrove, Richard C.; et al.

JOURNAL

Sport Performance & Science Reports

DATE DEPOSITED

6 April 2018

This version available at

<https://research.stmarys.ac.uk/id/eprint/2191/>

COPYRIGHT AND REUSE

Open Research Archive makes this work available, in accordance with publisher policies, for research purposes.

VERSIONS

The version presented here may differ from the published version. For citation purposes, please consult the published version for pagination, volume/issue and date of publication.

Impact of Fixture Congestion on Indices of Performance & Recovery in Youth Soccer Players

Chris Hattersley¹, Carl Wells², Richard Blagrove³, Steven Trangmar⁴, Stephen Patterson¹

¹School of Sport, Health & Applied Science, St Mary's University, London, UK, ²Sheffield United FC, UK, ³Faculty of Health, Education and Life Sciences, School of Health Sciences, Birmingham, UK, and ⁴Department of Life Sciences, University of Roehampton, Whitelands College, London, UK

Soccer | Fatigue | Recovery | Match congestion

Headline

Elite soccer schedules often require players to undertake two competitive fixtures within seven days. Previous research has reported that around 72-96 hours are required after a single soccer match for measures of muscle function to return to pre-game levels (1,2). As players are regularly required to complete periods where games are separated by only 48 to 72 hours contemporary research has started to investigate the effects fixture congestion has on elite senior players activity profiles and markers of recovery (3,4,5). However, it is apparent that there is a paucity of literature regarding the impact of fixture congestion in elite youth players (20).

Aim. It is unclear how the exposure to two competitive fixtures within seven days affects physical performance and recovery status in elite youth soccer players. The current study sought to identify if a difference existed in indices of physical output and recovery during the second fixture in a double game week in comparison to a single game week.

Methods

Athletes. 9 elite male youth soccer players (mean age 17.6 ± 0.6 years, stature 177.1 ± 6.9 cm, body mass 71.48 ± 6.9 kg) participated in two single game weeks and two double game weeks. All matches analysed during the study were competitive home fixtures from the English Under-18 Professional Development League 2 Northern Division. Ethical approval was granted by St Mary's University. Organisational, parent and participant consent was also obtained.

Design. A within group repeated measures design was used to examine the physical output and markers of recovery. To be included in the study each participant had to meet the following criteria (i) the player completed the full duration of all games (ii) the player didn't suffer any type of injury during any of the games (iii) the player played in the same position during each game. In addition, to control contextual factors (3), the following criteria had to be met in order for data to be included in the study (i) the first and second half of the game had no more than 3 added minutes of playing time (ii) the winning score line did not differ by more than 2 goals (iii) all games were played on the same pitch. Single game week fixtures were defined as the only competitive fixture occurring during a normal training week, with the game played on a Saturday. A double game week fixture was defined as the second fixture occurring during a week with two competitive games played on a Saturday – Tuesday schedule. Single game week and double game week fixtures took place over alternate weeks during the second half of the season. Training load was kept consistent throughout and recovery practices were standardised.

Methodology. To measure physical performance during the game a 10Hz GPS system (OptimEye X4, Catapult Innova-

tions, Canberra, ACT, Australia) was used to conduct time-motion analysis of the total distance (m), high intensity distance ($\geq 5.5\text{ms}^{-1}$) and amount of high intensity accelerations ($>3\text{ms}^{-2}$). Heart rate analysis (Polar Team2, Polar Electro Ltd, Warwick, England) was used to monitor cardiovascular exertion while an RPE load (RPE using the Borg CR10 scale x game time) was measured to quantify psycho-physiological fatigue (2,6). Indices of recovery were collected using a validated 0-100mm visual analogue scale (VAS) (7,8) for perceived recovery status, leg soreness, and sleep quality. In addition to this, a countermovement jump assessment was also completed on an integrated dual portable force plate set up with a sampling rate of 1000Hz (Pasco 2-axis force platform, Pasco, Roseville, CA, USA). Recovery measures were obtained at 1, 24, 48 and 72 hours post-game for all fixtures. Participants were familiarised with all data collection procedures during the 3 previous home fixtures that took place before the study.




Fig. 1. Comparison of perceived leg soreness VAS scores (mean ± SD) after SGW and DGW fixtures at 1 hour, 24 hours, 48 hours and 72 hours post game. *Small effect size, **moderate effect size, ***large effect size.




Fig. 2. Comparison of perceived recovery status VAS scores mean ± SD after SGW and DGW fixtures at 1 hour, 24 hours, 48 hours and 72 hours post game. *Small effect size, **moderate effect size, ***large effect size.

Table 1. Mean ± standard deviation of Visual Analogue Scale (VAS) measurements for perceived leg soreness and perceived recovery at 1, 2, 48 and 72 hours post-game.

Parameter	Time of Measurement Post - Game	Single Game Week	Double Game Week	Effect Size	95% CI Between Means	Qualitative
Perceived leg Soreness (VAS m)	1 hour	23.7 ± 10.4	18.5 ± 9.9	-0.51	-11, 1	Likely moderate decrease
	24 hours	47.9 ± 10.8	31.8 ± 9.3	-1.58	-22, -11	Likely large decrease
	48 hours	70.1 ± 8.0	43.0 ± 9.4	-3.08	-35, -19	Most likely large decrease
	72 hours	87.3 ± 5.4	72.3 ± 6.6	-2.46	-21, -8	Very likely large decrease
Perceived recovery Status (VAS mm)	1 hour	21.7 ± 9.4	14.8 ± 9.7	-0.71	-11, -2.5	Possible moderate decrease
	24 hours	44.2 ± 10.2	31.9 ± 10.8	-1.17	-17, -7.5	Very likely large decrease
	48 hours	70.8 ± 6.8	45.0 ± 7.7	-3.53	-31, -21	Most likely large decrease
	72 hours	96.0 ± 5.5	70.8 ± 7.4	-3.80	-24, -15	Very likely large decrease

Effect sizes (ES), 95% confidence intervals (CI) and magnitude based inferences (MBI) are provided between all measurements between single and double game weeks.

Table 2. Mean ± standard deviation of total distance (TD), high intensity distance (HID), high intensity accelerations (HIA), heart rate minutes over 85% maximum MHR (HR>85%) and rating of perceived exertion load (RPE Load).

Parameter	Game period	Single Game Week	Double Game Week	Effect Size	95% CI Between Means	Qualitative
Total Distance (m)	First Half	5734 ± 234	5592 ± 181	-0.67	-320, 38	Possible moderate decrease
	Second Half	5523 ± 226	5390 ± 223	-0.59	-290, 27	Possible moderate decrease
	Total	11257 ± 429	10982 ± 353	-0.69	-500, -46	Possible moderate decrease
High Intensity Distance (m) (≥5m.s-1)	First Half	401 ± 121	376 ± 100	-0.22	-63, 13	Possible small decrease
	Second Half	380 ± 133	297 ± 88	-0.73	-130, -35	Likely moderate decrease
	Total	781 ± 249	673 ± 186	-0.49	-180, -36	Very likely small decrease
High Intensity Accelerations (n) (>3m.s-2)	First Half	145 ± 24	134 ± 28	-0.42	-24, 0	Likely small decrease
	Second Half	135 ± 19	129 ± 23	-0.26	-118, -6	Possible small decrease
	Total	280 ± 42	263 ± 51	-0.36	-36, 5	Likely small decrease
Heart Rate >85% MHR (mins)	First Half	32 ± 3	31 ± 2	-0.43	-4, 0	Likely small decrease
	Second Half	31 ± 2	32 ± 3	0.35	-2, 3	Unclear
	Total	64 ± 4	63 ± 5	-0.07	-5, 3	Unclear

Effect sizes (ES), 95% confidence intervals (CI) and magnitude based inferences (MBI) are provided between first half, second half and total measures for all parameters between single and double game weeks.

Statistical analysis. 9 players met the eligibility criteria to be included in the study. All data are provided as means ± standard deviations. In line with the suggestions of Carling et al (3), Cohens d effect sizes (ES) and 95% confidence intervals (CI) were calculated to determine the magnitude of the differences between parameters from the single game and double game weeks. ES was classified as trivial (<0.2), small (0.2-0.6), moderate (0.6-1.2) and large (>1.2). A published spreadsheet (17) was used to make a qualitative probabilistic mechanistic inference about the true effect. Threshold values were calculated by dividing the between-subject standard deviation by 0.2 (small), 0.6 (moderate) or 1.2 (large) in order to determine the magnitude band that the effect falls into (18). The resulting values were translated into descriptors using the modified thresholds proposed by Batterham and Hopkins (19): 0-0.5% most unlikely; 0.5-5% very unlikely; 5-25% unlikely; 25-75% possibly; 75-95% likely; 95-99.5% very likely; and >99.5% most likely. If the probabilities of the effect being substantially positive and negative were both >5%, the effect was reported as unclear. Pearson correlations were used to investigate the relationship between physical performance and recovery data.

Results

The greatest difference between double game and single game weeks was highlighted by perceived measures of recovery status and muscle soreness which displayed large negative effect sizes after 24, 48 and 72 hours (Figure 1/2, Table 1). No

meaningful differences were found for CMJ or perceived sleep quality between single game and double game weeks at any time point (Due to this, no data for these parameters are included in the report but all data is included in the accompanying spreadsheet). There was a consistent pattern of physical output being reduced during a double game week when compared to a single game week (Table 2). Total distance and high intensity distance displayed the largest reductions during double game weeks with moderate negative effect sizes observed for both of these parameters.

Discussion

These findings are similar to those reported by Buchheit et al (20) who found that the majority of match running parameters were reduced for adolescents who were post peak height velocity during a period of congestion. The decrease in high speed running by 22% during the second half of double game week fixtures may have the biggest impact on performance as high-intensity actions are often the decisive factor involved in the outcome of a game (9). Furthermore, correlational analysis identified the strongest association between physical performance and markers of fatigue occurred between high intensity distance and post-match leg soreness (r= -0.81) highlighting that this may be the most fatiguing action. Previous studies have shown that high intensity running depletes glycogen stores (10) as well as causing more structural damage to the muscle (11), and an accumulation of these effects may have caused the decline in match running performance. In addition,

tion, Bradley and Noakes (12) reported that senior players adopt a self-pacing strategy in order to sustain high intensity running performance during periods of fixture congestion and the younger players in this study may not have developed this ability yet.

As indices of recovery were accentuated following double game week fixtures (Figure 1, Table 2) this suggests a residual fatigue from the previous fixture in that week might have contributed to the increased perception of fatigue following the second fixture. The magnitude of difference in fatigue effects was observed further away from match day and this is consistent with previous research that has shown delayed on-set muscle soreness (DOMS) peaks at 48-72 hours post-game (13). The findings are in agreement with the research carried out by Thorpe et al (14) that found subjective assessments are effective measurements of in-season fatigue in soccer players.

The overall findings of the study suggest that double game week fixtures may be more physically demanding than single game week fixtures for youth soccer players. The reduction in physical capacity and decline in indices of recovery were most likely caused by a transient state of fatigue characterised by muscular damage, glycogen depletion as well as an elevated amount of stress placed on the cardiovascular, endocrine, and nervous systems (11,15,16). This is likely attributable to the lower training age of youth players, meaning that they haven't yet developed the physical capacity and self-pacing strategies to consistently sustain match running performance levels. Additionally, the lower level of physical maturity may contribute to the fatigue effects experienced by young players being amplified.

Practical Applications

- An individual approach to player monitoring should be used, ensuring that a player's physical exertion and markers of recovery from the previous game are used to inform training load and use of appropriate recovery strategies.
- Despite the heightened levels of fatigue after double game weeks, young players may need to periodically be exposed to periods of congestion in order to develop the self-pacing strategies utilised by senior players. However, further research is needed to determine the association between fixture congestion and injury risk for youth players.
- Elite youth players should follow a long term development plan in order to improve their physical capacity and make them more resistant to fatigue. This should include sufficient exposure to match specific actions such as high speed running, accelerations and decelerations.

Limitations

- As the study was carried out in the real world setting during competitive fixtures it is difficult to meet the necessary inclusion criteria across a greater amount of games. Due to this, further research is required pooling data from multiple teams across a larger number of games.
- Pre-game measures could have been included for the markers of recovery (CMJ & VAS) to quantify the pre / post game fatigue effect as well as the time course of recovery rates post game.
- Although the main purpose of this study was to identify the general effects of fixture congestion for youth players, further analysis could be carried out to monitor these effects for specific playing positions.

Twitter: Follow Chris Hattersley @hattersley4, Carl Wells @CWsports, Richard Blagrove @rich.blagrove, Steven Trangmar @StevenTrangmar and Stephen Patterson @stephen_patt

References

1. Asencio A, Rebelo A, Oliveira E, Marques F, Periera L, Magalhaes J. Biochemical impact of a soccer match - analysis of oxidative stress and muscle damage markers throughout recovery. *Clin Biochem.* 2008 Jul;41(10):841-851.
2. Nedelec M, Mccall A, Carling C, Legall F, Berthoin S, Dupont G. Recovery in soccer: Part 1 - post match fatigue and time course of recovery. *Sports Med.* 2012 Dec;42(12):997-1015.
3. Carling C, Gregson W, Mccall A, Moreira A, Wong P, Bradley P. Match running performance during fixture congestion in elite soccer: Research issues and future directions. *Sports Med.* 2015 May;45(5) 10:605-613.
4. Dellal A, Lago-Penas C, Rey E, Chamari K, Orhant E. The effects of a congested fixture schedule on physical performance, technical activity and injury rate during matches in a professional soccer team. *Brit J Sports Med.* 2013 Feb;49(6):1-5.
5. Djaoui L, Wong D, Pialoux V, Hautier C, Da Silva C, Chamari K, Dellal A. Physical activity during a prolonged congested fixture period in a top-class European football team. *Asian J Sports Med* 2014 Mar;5(1):47-53.
6. Rampinini E, Bosio A, Ferraresi A, Petruolo A, Morelli A. Match-related fatigue in soccer players. *Med Sci Sport Exer.* 2011 Nov;43(11):2161-2170.
7. Morgan P, Brown R, Raglin S, O'Connor J, Ellickson A. Psychological monitoring of overtraining and staleness. *Br J Sports Med.* 1987 Sep;21(3):107-114.
8. Raglin S, Morgan P. Development of a scale for use in monitoring training-induced distress in athletes. *Int J Sports Med.* 1994 Feb;15(2):84-88.
9. Faude O, Koch T, Meyer T. Straight sprinting is the most frequent action in goal situations in professional football. *Journal of Sports Sciences.* 2012 Mar;30(7):625-631.
10. Krstrup P, Ortenblad N, Nielsen J, Nybo L, Gunnarson P. Maximal voluntary contraction force, function and glycogen resynthesis during the first 72 h after a high-level competitive soccer game. *Eur J App Physiol.* 2011 Dec;111(12):2987-2995.
11. Thorpe R, Sunderland C. Muscle damage, endocrine and immune marker response to a soccer match. *J Strength Cond Res.* 2011 Oct;26(10):2783-2790.
12. Bradley P, Noakes T. Match running performance fluctuations in elite soccer: Indicative of fatigue, pacing or situational influences? *J Sport Sci.* 2013 Jul;31(15):1627-1638.
13. Ispirlidis, I., Fatouros, I., Jamurtas, A., Nikolaidis, M., & Michalis, G. Time-course of changes in inflammatory and performance responses following a soccer game. *Clin J Sport Med.* 2008 Sep;18(5):423-431.
14. Thorpe R, Strudwick A, Buchheit M, Atkinson G, Drust B, Gregson W. Monitoring Fatigue During the In-Season Competitive Phase in Elite Soccer Players. *Int J Sports Phys Perf.* 2015 Feb;10(8):958-964.
15. Bangsbo J, Laia F, Krstrup P. Metabolic response and fatigue in soccer. *Int J Sports Physiol Perform.* 2007 Jun;2(2):111-127.
16. Krstrup P, Mohr M, Steensberg A, Bencke J, Kjaer M, Bangsbo J. Muscle and blood metabolites during a soccer game: Implications for sprint performance. *Med Sci Sport Exer.* 2006 Jun;38(6):1165-1174.
17. Hopkins WG. A spreadsheet for deriving a confidence interval, mechanistic inference and clinical inference from a P value. *Sportscience.* 2007 Jan 1;11:16-21.

18. Hopkins WG. Linear models and effect magnitudes for research, clinical and practical applications. *Sports Science*. 2010 Jan 1;14(1):49-57.
19. Batterham AM, Hopkins WG. Making meaningful inferences about magnitudes. *International journal of sports physiology and performance*. 2006 Mar;1(1):50-7.
20. Buchheit M, Horobeanu C, Mendez-Villanueva A, Simpson BM, Bourdon PC. Effects of age and spa treatment on match running performance over two consecutive games in highly trained young soccer players. *Journal of sports sciences*. 2011 Mar 1;29(6):591-8.

Copyright: The articles published on *Science Performance and Science Reports* are distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.