

Association of Daily Workload, Wellness, Injury and Illness during Tours in International Cricketers

¹ Robert Ahmun, ^{1,2,7}Steve McCaig, ³Jamie Tallent, ⁴Sean Williams, ^{5,6}Tim Gabbett,

¹England and Wales Cricket Board, Leicestershire, UK

²English Institute of Sport, Loughborough University, Loughborough,

³School of Sport Health and Applied Science, St Mary's University, Twickenham, UK

⁴Department for Health, University of Bath, Bath, UK

⁵Gabbett Performance Solutions, Brisbane, Australia

⁶Institute for Resilient Regions, University of Southern Queensland, Ipswich, Australia

⁷ School of Sports Science, Cardiff Metropolitan University, Cardiff, UK

Running Head: Workload and Illness associated with Touring Teams

Submission Type: Original Investigation

Address for Correspondence:

National Cricket Performance Centre

Loughborough University

Leicestershire

UK

LE11 3TU

Tel: 01509 228664

Fax: 01509 453718

Email: rob.ahmun@ecb.co.uk

ABSTRACT

Purpose: To examine the relationship between player internal workloads, daily wellness monitoring, injury and illness in a group of elite adolescent cricketers during overseas competitions. **Methods:** Thirty-nine male international adolescent cricketers (17.5 ± 0.8 yr) took part in the study. Data was collected over five tours across a three-year period (2014-2016). Measures of wellness were recorded, and daily training loads calculated using session-rating of perceived exertion. The injury and illness status of each member of the squad was recorded daily. Acute and chronic workloads were calculated using three days and fourteen days moving averages. Acute workloads, chronic workloads, and acute chronic workload ratios (ACWR) were independently modelled as fixed effects predictor variables. **Results:** In the subsequent week, a high 3-day workload was significantly associated with an increased risk of injury (Relative Risk [RR] = 2.51; CI = 1.70 to 3.70). Similarly, a high 14-day workload was also associated with an increased risk of injury (RR = 1.48; CI = 1.01 to 2.70). Individual differences in the load injury relationship were also found. No clear relationship between the ACWR and injury risk was found, but high chronic workloads combined with a high or low ACWR showed an increase probability of injury compared to moderate chronic workloads. There were also trends for sleep quality and cold symptoms worsening the week before an injury occurred. **Conclusion:** Although there is significant individual variation, short term high workloads and changing in wellness status appear to be associated with injury risk.

Key Words: Cricket, Workloads, Injury, Wellness

62

63 INTRODUCTION

64 It is well established that injury rates can influence the success of a team ¹ and consequently
65 managing loads appear to be an essential part of reducing injury risk. Training loads
66 comprise of both internal and external loads. External load relates to the amount of work
67 completed, whilst internal loads are a measure of the relative physiological strain. This
68 relationship is crucial in determining the stress and adaptive response ². Furthermore, the
69 rate of loading is a critical factor in influencing performance and injury factor ³. If loads are
70 applied in a moderate and progressive manner, they may be protective against injury ². No
71 single marker can be used to accurately predict when an athlete enters a maladaptive state,
72 so a combination of both internal and external load measures, specific to the nature of the
73 sport, is recommended ^{3,4}.

74

75 Despite the increased use of global positioning system (GPS) to record load in the literature
76 ⁵, external load of cricket fast bowling is predominantly measured using the number of balls
77 bowled ^{6,7}. However, recently balls bowled has been shown to inadequately capture the cost
78 of fast bowling ⁸. Consequently a hybrid of the session-rating of perceived exertion (sRPE) ⁹,
79 TRIMP and balls bowled has been used to model injury risk in cricket. Hulin, Gabbett,
80 Blanch, Chapman, Bailey, Orchard ¹⁰ was the first to investigate this specifically in cricket
81 and combined both external (balls bowled) and internal load data (sRPE x duration) to
82 model injury risk in fast bowlers. Despite a significant relationship between acute (1-week)
83 external workloads and increased injury risk in the current week, no relationships were
84 demonstrated between sRPE's and injury risk in the current or subsequent week. However,
85 when both the external and internal acute workload exceeded chronic (4-week rolling
86 average) workload, resulting in an acute chronic workload ratio (ACWR) of >2.0, the relative
87 risk of fast bowling injury was 3.3 to 4.5 times greater. As balls bowled does not appear to
88 accurately reflect external workload ⁸, internal workloads may be more strongly associated
89 with injury as it encompasses all aspects of training and competition.

90

91 It is also highlighted that progressively higher workloads may serve as a protective
92 mechanism against injury ¹⁰. Unfortunately, progressive sequenced training to develop high
93 chronic training loads is not always feasible, particularly in adolescent cricket where

overseas tours occur out of season. The nature of touring results in intensive training periods followed by a congested fixture period. These, intensive training periods have led to an increase catabolic environment during the competition period ¹¹. Whilst an increased catabolic environment does not necessarily directly influence performance, it can indicate the ability of the athletes to tolerate training load ¹². Short duration tours have resulted in an increased injury risk in many other sports ^{13,14} although it is unclear if cricket has similar traits. Even though a significant amount of a cricketer's career is spent touring various countries, the effect this has on injury risk is unknown. As less recovery between days of bowling has been shown to increase the risk of injury in young (14.7 ± 1.4 years) fast bowlers ⁶, it is hypothesised that touring would also be associated with a high risk of injury. A recent systematic review has highlighted the large quantity of self-reported measures of wellness that are used in sport ¹⁵. However, despite this review the relationship with injury and well-being is inconclusive. Therefore, the aim of this study was to examine the relationship between internal workloads, daily wellness scores, injury and illness in a group of elite adolescent cricketers during overseas competitions.

METHODS

Participants

The sample comprised 39 male adolescent cricketers (17.5 ± 0.8 y) who were selected to play international age group cricket. Data were collected over five tours across a three-year period (2014-2016). Tour duration varied from 18 to 30 days with a mean tour duration of 24 ± 5 days. Of the five tours, 26% of the participants ($n = 10$) played one tour, 53% ($n = 20$) played two tours and 21% ($n = 8$) played three tours – equating to 1862 training days. Data were collected as a part of the routine practices throughout the tour season to which all players had consented ¹⁶. The project was approved by St Mary's University Ethics Committee in the spirit of the Helsinki Declaration.

Wellness Data

Subjective measures of wellness were recorded each morning at breakfast using a five-point scale for sleep quality and duration, muscle soreness, cold symptoms and mood with lower scores being indicative of reduced wellness ¹⁷.

126 Internal Workload

127 Players were asked to provide a subjective rating of perceived exertion (RPE) using a 10-
128 point rating scale ⁹. The intensity of all sessions (games, cricket training and strength and
129 conditioning) were recorded within 30 mins of completing the session. Daily training loads
130 were then calculated by multiplying session RPE by session duration (min).

131

132 Injury Data Collection

133 The programme's physiotherapist collected the data throughout the course of the study and
134 the same practitioner was the programme physiotherapist for the duration of the study. A
135 programme day was defined as any day where the squad was together, be it for a match,
136 training, rest or travel day. For each programme day the squad physiotherapist recorded the
137 injury status of each member of the squad on a specifically designed spreadsheet. A broader
138 definition of injury and illness, as used in the current study, provides a more complete
139 picture of the true burden of injury and illness than a time loss definition of injury and
140 illness. The recent international injury consensus statement on injury surveillance¹⁸ in
141 cricket updated its definition of a cricket injury to include medical attention conditions and
142 our paper is consistent with this consensus statement. Each player's injury status was
143 recorded as being either:

144 1. Fully available for training and matches, with no injury or illness

145 2. Fully available for training and matches, but with an injury or illness

146 3. Available for selection in a major match, but with modified activity due to injury or
147 illness

148 4. Unavailable for selection in a major match due to injury or illness

149 Non time-loss injuries were category 2 and 3 and time-loss injuries were category 4. All new
150 injuries, as well as any pre-existing injuries players carried into the programme were
151 reported. It was possible for a player to have multiple injuries or illness at any one time e.g.

they may have a medical condition while being treated for a musculoskeletal condition or they may have two or more musculoskeletal conditions requiring management at the same time.

A change in injury status occurred when a player's injury status changed from one to another e.g. a player sustained a hamstring strain and was unavailable for selection (category 4), but the previous day they were fully available with no injury or illness (category 1). Only injury status changes where the players' condition worsened i.e. they required increasing medical attention or activity/participation restriction; were included for analysis. This was a negative injury status change. This occurred when their injury status category number increased and was considered a negative status change.

For each injury or illness, the squad physiotherapist also recorded the players skill group, the side, region and location of injury, diagnosis based on the Orchard sports injury classification system 10 (OSICS10) ¹⁹ and the number of programme days spent in each injury status category. In addition, the mode of injury onset, activity at the time of onset and whether it occurred on a match or non-match day was recorded as well. Skill group was defined as per the international consensus statement guidelines ¹⁸, with players classed as either batsman, pace bowlers, slow bowlers or wicketkeepers. The mode of onset followed the consensus statement guidelines (Orchard, Ranson, Olivier et al, 2016), and was defined as either sudden onset, impact (blow or contact), gradual onset, insidious or illness. Sudden onset injuries comprised non-impact muscle strains and ligament sprains e.g. an ulnar collateral ligament sprain during a one-off throw. Impact injuries occurred because of contact with another player or object e.g. a contusion due to being hit by the ball. A gradual onset injury was where the condition developed over time e.g. a rotator cuff tendinopathy from repetitive throwing. An insidious onset was where there was no identifiable activity associated with a musculo-skeletal injury. Illness was any medical condition not associated with the other four mechanisms.

180

181 Data Analysis

182 'Programme' exposure was calculated by multiplying the number of players in each squad
183 during each day of the programme by the number of programme days, using the following
184 formula:

$$185 \quad \textit{Programme player days} = (\textit{daily squad size} \times \textit{number of days on programme})$$

186

187 Statistical Analysis

188 All estimations were made using the lme4 package with R (version 3.3.1, R Foundation for
189 Statistical Computing, Vienna, Austria). Acute and chronic workloads were calculated using
190 exponentially-weighted moving averages with time constants of three days and fourteen
191 days, respectively ²⁰. These time frames were chosen to reflect the 'tour' format of the
192 competitions analyzed (i.e., 18 to 30 day tours with limited chronic loading) and because
193 exponentially-weighted moving averages have shown stronger relationships with injury risk
194 than the usual one and four week rolling periods ²¹. Uncoupled ACWR were calculated by
195 reporting acute workloads (i.e., fatigue) as a proportion of chronic workloads (i.e., fitness)
196 ¹⁰, such that acute load periods were not included in the calculation of chronic load ²².
197 Within-individual Z-scores were calculated for each player using the following formula:

198 $(\text{individual player's score} - \text{individual player's average}) / \text{individual player's standard}$
199 deviation ; a Z-score is the number of standard deviations the response is above or below
200 the mean of the distribution.

201 A generalized linear mixed-effects model (GLMM) was used to model the association
202 between workloads and injury risk in the subsequent week. This mixed effects model was
203 selected for its ability to account for repeated measurements, and to explore individual
204 responses between workloads and injury risk. Acute workloads, chronic workloads, and
205 ACWR were independently modelled as fixed effects predictor variables. In addition, the

interaction between ACWR and both acute and chronic workloads was assessed by including multiplicative terms in the model. Random effects were athlete identity (differences between athletes' mean injury risk), athlete \times tour (variability within athletes between tours), and the residual. If assessment of a quadratic trend between the workload measure and injury risk was significant ($P \leq 0.05$), the measure was split into tertiles for analysis, with the lowest load range being the reference group. Otherwise, linear effects for continuous predictor variables were evaluated as the change in relative injury risk (RR) associated with a two standard deviation increase in the workload or wellness measure (representing the change associated with a 'typically low' versus a 'typically high' value of the predictor)²³. The odds ratios obtained from the GLMM model were therefore converted to RR in order to interpret their magnitude²⁴. The RR represents the change in injury risk associated with changes in the investigated load or wellness variables. A RR of 1.0 represents no change in risk of injury, whilst values of 0.5 and 2.0 would represent a halving or doubling of injury risk, respectively.

Magnitude-based inferences were used to provide an interpretation of the real-world relevance of the outcomes²⁵. The smallest important increase in injury risk was a RR of 1.11, and the smallest important decrease in risk was 0.90²⁵. An effect was deemed unclear if the chance that the true value was beneficial was $>25\%$, with odds of benefit relative to odds of harm (odds ratio) of <66 . Otherwise, the effect was deemed clear, and was qualified with a probabilistic term using the following scale: $<0.5\%$, most unlikely; $0.5-5\%$, very unlikely; $5-25\%$, unlikely; $25-75\%$, possible; $75-95\%$, likely; $95-99.5\%$, very likely; $>99.5\%$, most likely²⁶. The *r2glmm* package was used to determine whether model fit was significantly improved when using GLMM in comparison with a logistic regression model (which does not account for repeated measurements or individual variations in responses).

RESULTS

Thirty-nine players were involved in 1862 programme days during the study. There were 98 injuries in 38 players that resulted in 130 negative injury status changes on 125 different programme days. Only 17 (13.1%) of these changes resulted in the player being unavailable for match selection (category 4). On average players had a negative injury status change every 14.3 days. In most negative status changes (53.1%) players went from being fully

available to receiving medical attention (change status from category 1 to 2), the next most common status changes were from fully available to modified activity (category 1 to 3) with 20% of all changes; and from medical attention to modified activity (category 2 to 3) with 13.8% of all changes. Sixteen pace bowlers accounted for 43.7% of all programme days and 46.9% of all negative status changes, nine spin bowlers accounted for 24.9% of programme days and 26.9% of all changes, nine batsmen accounted for 19.8% of programme days and 15.4% of all changes and five wicketkeepers accounted for 11.6% of programme days and 10.8% of all changes. Compared to pace bowlers (RR = 1.00 (ref)), wicket keepers and batsmen had a lower overall risk of injury (RR = 0.56; CI = 0.29 to 1.08), whilst the inference for spin bowlers was unclear (RR: 0.70, CI = 0.31 to 1.57).

****Insert Table 1 here****

Wellness and Injury Risk

No relationship was found between wellness scores and injury risk in the subsequent week, although there were trends for sleep quality and cold symptoms to worsen the week before an injury occurred (Table 1).

****Insert Table 2 here****

Acute and Chronic Workloads

In the subsequent week, a high (>0.35) 3-day workload z-score was significantly associated with an increased risk of injury (RR = 2.51; CI = 1.70 to 3.70; likelihood range >99.5%, most likely), compared with medium (-0.45 to 0.35) and low (<-0.45) workload z-scores (Table 2). The predicted probability of injury increased from 6% to 11% as 3-day workload increased from medium to high categories. This is in comparison to overall risk of pace bowlers (RR = 1.00 (ref)), wicket keepers and batsmen (RR = 0.56; CI = 0.29 to 1.08), spin bowlers (RR: 0.70, CI = 0.31 to 1.57).

****Insert Table 3 here****

A high (>0.67) 14-day workload z-score was also associated with an increased risk of injury (RR = 1.48; CI = 1.01 to 2.70; likelihood range 75-95%, likely), compared with medium (-0.45 to 0.35) and low (<-0.45) workload z-scores (Table 3). The predicted probability of injury increased from 8% to 13% as 14-day workload increased from medium to high categories.

****Insert Table 4 here****

The ACWR was not clearly associated with injury risk (Table 4). Both acute and chronic workloads were independently associated with injury risk in a linear fashion (Figure 1), with 2 standard deviation increases in both predictors (620 AU and 538 AU, respectively) associated with substantial increases in injury risk (Acute: RR: 1.82, CI = 1.34 – 2.47, most likely harmful; Chronic: RR: 2.22, CI: 1.56 – 3.15, most likely harmful).

****Insert Figure 1 here****

Additionally, there was a clear interaction effect between ACWR categories and chronic workloads (Figure 2), such that the effect of increasing chronic workloads on injury risk was substantially higher in the 'low' and 'high' ACWR categories, compared to the 'moderate' ACWR category. There was no interaction effect observed between ACWR and acute workloads ($P = 0.30$).

****Insert Figure 2 here****

There was a substantial improvement in model fit when random effects were included in the model (logistic regression model $R^2 = 10\%$, GLMM model $R^2 = 27\%$, $P < 0.001$). Therefore, individual differences in workload-injury relationships were evident. Figure 3 displays the relationship between chronic workloads and injury risk in the subsequent week for each individual in the analysis, as estimated via the GLMM.

****Insert Figure 3 here****

DISCUSSION

This is the first study to establish specific workload thresholds for adolescent cricketers and also non-fast bowlers. The study had numerous key findings. Firstly, high short-term (3 days) workloads (>2125 AU) or a high 14 day workload (>7212 AU) were also associated with an increased injury risk. Secondly, high chronic loads combined with a high or low ACWR increases the probability of injury compared to moderate chronic loads. Thirdly, individual differences in injury risk was also demonstrated between players. Finally, sleep quality and cold symptoms showed a trend with injury risk.

The findings from our study show that high short-term workloads in cricket (>2125 AU) increase the risk of injury. High workloads and increased injury risk may be a result of inadequate recovery time between sessions. Particularly during the early days of touring, an optimal balance between intensity and volume of training and recovery needs to be implemented. These findings are in-line with previous sports such as rugby²⁷ and football²⁸. In contrast to our findings, previous work in cricket¹⁰ has found no link between acute internal workload measures and injury risk. One explanation for the difference could be the age and experience of the players involved in the study. Hulin, Gabbett, Blanch, Chapman, Bailey, Orchard¹⁰ used older (26 ± 5 yr), more experienced cricketers who have had exposure to higher chronic workloads compared to the younger (17.5 ± 0.8 years) adolescent cricketers in our study. Individuals with a greater physical training history or greater physical attributes have also shown better tolerance to acute spikes in load better than younger individuals²⁹. Therefore, the finding of greater injury risk with rapid acute changes in load in adolescent cricketers may be expected. Conversely, older athletes appear to be at greater risk of injuries at a given absolute training load than younger athletes³⁰. Whilst this appears to be a contradictory finding, there may be a 'sweet spot' for age, physical qualities and training history where athletes can cope with acute spikes in training loads. Other differences between the findings in this study may due to the classification or change in injury states we used compared to time loss data¹⁰. Our study also used a change in injury status that is more reflective of current sporting practices.

The non-significant relationship between the ACWR and risk of injury or illness is in contrast to previous work in senior cricket fast bowlers¹⁰ and elite adolescent cricketers³¹. Our

findings uniquely show that high chronic loads combined with a high or low ACWR increases the probability of injury compared to moderate chronic loads. Previously, higher chronic workloads have shown to serve as an injury protective mechanism for acute spikes in workload ³². Conversely, high chronic loads can be achieved safely so long as the ACWR is not excessive. Despite being beyond the scope of this study, it seems essential that the workload prior to touring is recorded. If players have accumulated large workloads before touring then ensuring ACWR is not minimised or excessive would appear to reduce the risk of injury.

Individual differences in injury risk were also demonstrated between playing positions for the first time showing that athletes should understand individual responses to chronic workloads. Prescribing individual load is often very difficult in a team setting, but our data suggests that ensuring all players are below (>2125 AU) will reduce the risk of injury. The length of the acute window has also been shown to be strongly associated with injury ³³. Given that players do not have the opportunity to build chronic workloads prior to touring, our study used time frames of 3 and 14 days for acute and chronic loading periods. Work has predominantly used time frames of 7 and 28 days though there is evidence to suggest that 6 and 21 days acute to chronic workload ratios is optimal for predicting injuries ³³. Consequently, it could be suggested that the 3 days used for the acute period in our study was not long enough to see differences in ACWR.

A positive link between alterations in training load and subjective measures of well-being has previously been established ^{34,35}. A recent systematic review ¹⁵ has highlighted that subjective well-being measures respond consistently to stress imposed by training. Of 56 research articles, 85% favoured subjective measures when monitoring athlete load. Negative changes in wellness measures have also been linked to increased risk of illness ³⁶, although changes in wellness measures and risk of injury has received less attention ³⁷. The result from our study showed no significant relationship between subjective measures and injury and illness. A possible explanation for these findings may be due to the scale used. Our study used a 5-point scale where previous work has shown that a greater number of points on a scale increases the sensitivity ³⁸. However, we did observe trends of reduced sleep and self-reported cold symptoms in the week before an injury occurred. Recent work

by von Rosen, Frohm, Kottorp, Friden, Heijne ³⁷ supports this notion and demonstrated that in youth athletes, an increase in training load and intensity in addition to a decrease in sleep volume significantly increased the risk of injury. With even modest sleep loss associated with impairment of psychomotor performance ³⁹ it appears logical that assessing sleep volume and quality is a key subjective measure for reducing injury and illness risk.

LIMITATIONS

Although higher chronic workloads have been shown to be associated with a lower risk of injury, it was not possible to quantify chronic training workloads in the period prior to tours. Therefore, future work should focus on the workloads preceding a tour and the effects this has on injury prevalence. Subjective measures of wellness were asked during breakfast. Whilst the experimenters made every attempt to ensure this was performed away from other coaches and players, the nature of the touring environment sometimes meant wellness measures were not performed in isolation. Finally, the nature of cricket often involves large periods of very low inactivity such as fielding in a match. This low RPE but long duration can often cause excessively large TRMP values.

CONCLUSION

Collectively, these results demonstrate that in elite adolescent cricketers, high acute and/or chronic internal workloads are significantly associated with an increased risk of injury. Rapid increases in acute workloads >2125 AU are more closely associated with injury and illness than ACWR assessed over 3 and 14 days respectively. High chronic loads combined with a high or low ACWR increases the probability of injury compared to moderate chronic loads. Therefore, practitioners should ensure individuals that accumulate large amounts of workload have a moderate ACWR whilst touring. However, the injury risk appears to be an individualised response. We have demonstrated for the first time that other cricket skill sets (in addition to fast bowling) have injury risks associated with workloads. Although not significant, measures of wellness, specifically sleep duration and self-reported cold symptoms can be expected to worsen the week before an injury occurs.

PRACTICAL IMPLICATIONS:

- 395 • The non-invasive and simple session-RPE method is useful for tracking training and
396 game loads and injury risk during elite adolescent cricket tours.
- 397 • Coaches should avoid spike in workloads when chronic workloads are high or low.
- 398 • Players appear to be at an increased risk of injury when they experience a high 3 day
399 cumulative load (≥ 2125 AU), though there are individual differences.
- 400 • Although not significant, worsening sleep quality and self-reported cold symptoms
401 are possible subjective indicators of heightened injury risk in this population. These
402 measures warrant further investigation in larger studies in the future.

403

404

REFERENCE LIST

1. Hagglund M, Walden M, Magnusson H, Kristenson K, Bengtsson H, Ekstrand J. Injuries affect team performance negatively in professional football: an 11-year follow-up of the UEFA Champions League injury study. *Br J Sports Med.* 2013;47(12):738-742.
2. Jones CM, Griffiths PC, Mellalieu SD. Training Load and Fatigue Marker Associations with Injury and Illness: A Systematic Review of Longitudinal Studies. *Sports Med.* 2017;47(5):943-974.
3. Soligard T, Schwellnus M, Alonso JM, et al. How much is too much? (Part 1) International Olympic Committee consensus statement on load in sport and risk of injury. *Br J Sports Med.* 2016;50(17):1030-1041.
4. Vickery W, Dascombe B, Duffield R. The Association Between Internal and External Measures of Training Load in Batsmen and Medium-Fast Bowlers During Net-Based Cricket Training. *Int J Sports Physiol Perform.* 2017;12(2):247-253.
5. Petersen CJ, Pyne D, Dawson B, Portus M, Kellett A. Movement patterns in cricket vary by both position and game format. *J Sports Sci.* 2010;28(1):45-52.
6. Dennis RJ, Finch CF, Farhart PJ. Is bowling workload a risk factor for injury to Australian junior cricket fast bowlers? *Br J Sports Med.* 2005;39(11):843-846; discussion 843-846.
7. Orchard JW, Blanch P, Paoloni J, et al. Cricket fast bowling workload patterns as risk factors for tendon, muscle, bone and joint injuries. *Br J Sports Med.* 2015;49(16):1064-1068.
8. Tallent J, Higgins M, Parker N, et al. Quantification of bowling workload and changes in cognitive function in elite fast bowlers in training compared with twenty20 cricket. *J Sports Med Phys Fitness.* 2017.
9. Foster C, Florhaug JA, Franklin J, et al. A new approach to monitoring exercise training. *J Strength Cond Res.* 2001;15(1):109-115.
10. Hulin BT, Gabbett TJ, Blanch P, Chapman P, Bailey D, Orchard JW. Spikes in acute workload are associated with increased injury risk in elite cricket fast bowlers. *Br J Sports Med.* 2014;48(8):708-712.
11. McNamara DJ, Gabbett TJ, Naughton G, Farhart P, Chapman P. Training and competition workloads and fatigue responses of elite junior cricket players. *Int J Sports Physiol Perform.* 2013;8(5):517-526.
12. Frx AC, Kraemer WJ, Stone MH, Koziris LP, Thrush JT, Fleck SJ. Relationships Between Serum Testosterone, Cortisol, and Weightlifting Performance. *Journal of Strength and Condition Research.* 2000;14(3):338-343.
13. Carling C, McCall A, Le Gall F, Dupont G. The impact of short periods of match congestion on injury risk and patterns in an elite football club. *Br J Sports Med.* 2016;50(12):764-768.
14. Dellal A, Lago-Penas C, Rey E, Chamari K, Orhant E. The effects of a congested fixture period on physical performance, technical activity and injury rate during matches in a professional soccer team. *Br J Sports Med.* 2015;49(6):390-394.
15. Saw AE, Main LC, Gustin PB. Monitoring the athlete training response: subjective self-reported measures trump commonly used objective measures: a systematic review. *Br J Sports Med.* 2016;50(5):281-291.

- 451 16. Winter EM, Maughan RJ. Requirements for ethics approvals. *J Sports Sci.*
452 2009;27(10):985.
- 453 17. McLean BD, Coutts AJ, Kelly V, McGuigan MR, Cormack SJ. Neuromuscular,
454 endocrine, and perceptual fatigue responses during different length between-match
455 microcycles in professional rugby league players. *Int J Sports Physiol Perform.*
456 2010;5(3):367-383.
- 457 18. Orchard JW, Ranson C, Olivier B, et al. International consensus statement on injury
458 surveillance in cricket: a 2016 update. *Br J Sports Med.* 2016;50(20):1245-1251.
- 459 19. Rae K, Orchard J. The Orchard Sports Injury Classification System (OSICS) version 10.
460 *Clin J Sport Med.* 2007;17(3):201-204.
- 461 20. Williams S, West S, Cross MJ, Stokes KA. Better way to determine the acute:chronic
462 workload ratio? *Br J Sports Med.* 2016.
- 463 21. Murray NB, Gabbett TJ, Townshend AD, Blanch P. Calculating acute:chronic workload
464 ratios using exponentially weighted moving averages provides a more sensitive
465 indicator of injury likelihood than rolling averages. *Br J Sports Med.* 2017;51(9):749-
466 754.
- 467 22. Lolli L, Batterham AM, Hawkins R, et al. Mathematical coupling causes spurious
468 correlation within the conventional acute-to-chronic workload ratio calculations. *Br J*
469 *Sports Med.* 2017.
- 470 23. Hopkins WG, Marshall SW, Batterham AM, Hanin J. Progressive statistics for studies
471 in sports medicine and exercise science. *Med Sci Sports Exerc.* 2009;41(1):3-13.
- 472 24. Hopkins WG, Marshall SW, Quarrie KL, Hume PA. Risk factors and risk statistics for
473 sports injuries. *Clin J Sport Med.* 2007;17(3):208-210.
- 474 25. Batterham AM, Hopkins WG. Making meaningful inferences about magnitudes. *Int J*
475 *Sports Physiol Perform.* 2006;1(1):50-57.
- 476 26. Hopkins WG. A spreadsheet for deriving a confidence interval, mechanistic inference
477 and clinical inference from a p value. *Sportscience.* 2007;11:16-20.
- 478 27. Cross MJ, Williams S, Trewartha G, Kemp SP, Stokes KA. The Influence of In-Season
479 Training Loads on Injury Risk in Professional Rugby Union. *Int J Sports Physiol*
480 *Perform.* 2016;11(3):350-355.
- 481 28. Malone JJ, Di Michele R, Morgans R, Burgess D, Morton JP, Drust B. Seasonal
482 training-load quantification in elite English premier league soccer players. *Int J Sports*
483 *Physiol Perform.* 2015;10(4):489-497.
- 484 29. Malone S, Hughes B, Doran DA, Collins K, Gabbett TJ. Can the workload-injury
485 relationship be moderated by improved strength, speed and repeated- sprint
486 qualities? *Journal of Science and Medicine in Sport.* 2018.
- 487 30. Rogalski B, Dawson B, Heasman J, Gabbett TJ. Training and game loads and injury risk
488 in elite Australian footballers. *J Sci Med Sport.* 2013;16(6):499-503.
- 489 31. Warren A, Williams S, McCaig S, Trewartha G. High acute:chronic workloads are
490 associated with injury in England & Wales Cricket Board Development Programme
491 fast bowlers. *J Sci Med Sport.* 2018;21(1):40-45.
- 492 32. Hulin BT, Gabbett TJ, Lawson DW, Caputi P, Sampson JA. The acute:chronic workload
493 ratio predicts injury: high chronic workload may decrease injury risk in elite rugby
494 league players. *Br J Sports Med.* 2015.
- 495 33. Carey DL, Blanch P, Ong KL, Crossley KM, Crow J, Morris ME. Training loads and
496 injury risk in Australian football-differing acute: chronic workload ratios influence
497 match injury risk. *Br J Sports Med.* 2017;51(16):1215-1220.

34. Coutts AJ, Wallace LK, Slattery KM. Monitoring changes in performance, physiology, biochemistry, and psychology during overreaching and recovery in triathletes. *Int J Sports Med*. 2007;28(2):125-134.
35. Coutts AJ, Reaburn P. Monitoring changes in rugby league players' perceived stress and recovery during intensified training. *Percept Mot Skills*. 2008;106(3):904-916.
36. Thornton HR, Delaney JA, Duthie GM, et al. Predicting Self-Reported Illness for Professional Team-Sport Athletes. *Int J Sports Physiol Perform*. 2016;11(4):543-550.
37. von Rosen P, Frohm A, Kottorp A, Friden C, Heijne A. Multiple factors explain injury risk in adolescent elite athletes: Applying a biopsychosocial perspective. *Scand J Med Sci Sports*. 2017;27(12):2059-2069.
38. Diefenbach MA, Weinstein ND, O'Reilly J. Scales for assessing perceptions of health hazard susceptibility. *Health Educ Res*. 1993;8(2):181-192.
39. Vgontzas AN, Zoumakis E, Bixler EO, et al. Adverse effects of modest sleep restriction on sleepiness, performance, and inflammatory cytokines. *J Clin Endocrinol Metab*. 2004;89(5):2119-2126.

Wellness	Relative risk	Lower CL	Upper CL	Inference	P-value
Total wellness	0.96	0.82	1.13	Unclear	0.70
Sleep duration	0.98	0.83	1.16	Unclear	0.87
Sleep quality	0.89	0.76	1.05	Possibly ↓	0.25
Body feeling	0.91	0.77	1.08	Possibly trivial	0.35
Cold symptoms	0.86	0.72	1.02	Possibly ↓	0.15
Mood	1.00	0.84	1.18	Unclear	0.98

Table 1. Change in injury risk associated with a 2SD improvement in self-reported wellness indicator.

574 **Table 2.** Acute and chronic workloads express as AU and z-scores

	Z-Score	sRPE AU's
Acute Workload (3 Days)		
Low	<-0.45	523 - 1322
Medium	-0.45 to 0.35	1323 - 2124
High	>0.35	>2125
Chronic Workload (14 Days)		
Low	<-0.40	2051 - 5128
Medium	-0.4 to 0.67	5129 - 7211
High	>0.67	>7212

575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595

596
597

598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616

Table 3. Predicated probability of injury expressed relative to z-scores.

	Relative risk	Lower CL	Upper CL	Inference	P-value
3-day load z-score					
Low (<-0.45)	1.00 (ref)				
Medium (-0.45 to 0.35)	1.18	0.73	1.93	Unclear	0.56
High (>0.35)	2.40	1.57	3.66	Most likely ↑	0.0007
14-day load z-score					
Low (<-0.40)	1.00 (ref)				
Medium (-0.40 to 0.67)	1.18	0.82	1.71	Unclear	0.46
High (>0.67)	1.89	1.26	2.85	Most likely ↑	0.01

617
618
619
620
621
622
623
624
625
626

Table 4. Predicted probability of injury risk.



Acute:chronic workload	Relative risk	Lower CL	Upper CL	Inference	P-value
Low (<0.80)	1.00 (ref)	-	-	-	-
Medium (0.80 to 1.30)	0.99	0.64	1.56	Unclear	0.99
High (>1.30)	1.01	0.65	1.58	Unclear	0.96

Figure Legends

Figure 1. Acute, chronic and acute chronic workload ratio probability of injury.

Figure 2. Acute, chronic and acute to chronic workload ratios association with injury risk.

Figure 3. Relationship between chronic workloads and injury risk in the subsequent week for each individual. Primary role is defined. BAT = Batsmant; PB = Pace Bowler; SP = Seam Bowler; WK = Wicket Keeper



Marginal effects of model predictors

