

Sex-based comparison of the blood pressure, haemodynamic and cardiac autonomic adaptations following isometric exercise training in sedentary adults: a randomised controlled trial

Harry Swift ^{1,2} Damian Coleman ² Charles Pedlar ^{1,3}
Neil Andrew Smart ⁴ Chris Farmer ⁵ Jonathan Wiles ²

To cite: Swift H, Coleman D, Pedlar C, *et al*. Sex-based comparison of the blood pressure, haemodynamic and cardiac autonomic adaptations following isometric exercise training in sedentary adults: a randomised controlled trial. *BMJ Open Sport & Exercise Medicine* 2025;**11**:e002431. doi:10.1136/bmjsem-2024-002431

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/bmjsem-2024-002431>).

Accepted 15 May 2025



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

For numbered affiliations see end of article.

Correspondence to

Dr Jonathan Wiles;
jim.wiles@canterbury.ac.uk

ABSTRACT

Objectives We aimed to explore sex-specific differences in resting blood pressure (BP) reduction and associated cardiovascular adaptations following isometric exercise training (IET).

Methods 100 sedentary adults with normal to high-normal systolic BP volunteered for the study. Participants either performed home-based lower-body IET in the form of a wall squat three times a week for 4 weeks (each session comprised 4×2 min bouts) or were allocated to the control group. Cardiovascular variables, including BP, cardiac output, total peripheral resistance, stroke volume and heart rate variability (HRV), were measured at rest preintervention and postintervention.

Results Following 4 weeks of IET, there were no significant differences in resting systolic BP and diastolic BP between females (122.1±6.9 and 80.0±8.3 mm Hg) and males (119.6±7.2 and 77.4±8.6 mm Hg). However, female participants had a greater cardiac autonomic response following training, evidenced by a lower low-frequency to high-frequency HRV ratio (F: 1.38±1.27 and M: 2.1±1.5, $p=0.004$) and decreased and elevated low-frequency normalised units (F: 50.3%±16.2% and M: 60.9%±16.9%, $p=0.015$) and high-frequency normalised units (F: 49.7%±16.2% and M: 39.1%±16.9%, $p=0.015$), respectively.

Conclusions While resting BP reductions were comparable between female and male participants, there was a greater autonomic response and a higher incidence of clinically important BP reductions in females, which could indicate a greater cardioprotective effect following IET. These findings highlight the importance of considering sex differences in the prescription and evaluation of exercise interventions for hypertension management.

INTRODUCTION

Cardiovascular disease (CVD) accounts for approximately 32% of all global deaths annually.¹ Hypertension (HTN) is a primary risk factor for CVD, as it increases the rates of atherosclerosis and other vascular changes that compromise cardiovascular function.²

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Isometric exercise training (IET) has effectively reduced resting blood pressure (BP), often outperforming other exercise modalities like aerobic and dynamic resistance training. While prior meta-analyses suggest no consistent sex differences in resting BP responses to IET, limited studies with small sample sizes have resulted in conflicting findings regarding sex-specific physiological adaptations.

WHAT THIS STUDY ADDS

⇒ This study is the largest primary investigation that directly compares sex-based resting BP reductions and physiological adaptations following IET, using a robust sample size and validated methodology. It demonstrates that while BP reductions were comparable between sexes, females showed a greater improvement in cardiac autonomic modulation, highlighting potential sex-specific mechanisms of cardiovascular adaptation.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ These findings call for further exploration of sex-specific mechanisms in exercise prescription, particularly in the context of autonomic regulation and hormonal influences. Sex-specific tailoring of exercise protocols could optimise cardiovascular outcomes, potentially influencing guidelines for hypertension management.

Recent data have demonstrated that age-standardised rates of HTN are similar between females and males (32% and 34%, respectively).³ Current 2023 European Society of Hypertension (ESH) guidelines recommend that females and males with suboptimal blood pressure (BP) and low cardiovascular risk engage in non-pharmaceutical lifestyle interventions such as daily physical activity.⁴ Notwithstanding this, the guidelines mention

that there may be differences in the magnitude of BP reduction between females and males when following the exercise recommendations presented. However, evidence for this statement is based on a single meta-analysis conducted over a decade ago that found a greater BP-lowering effect following exercise training in males following aerobic (AT) and dynamic resistance training (DRT),⁵ and there remains a lack of research in this area.

Recent data suggest that BP reductions following isometric exercise training (IET) are greater than following AT or DRT.⁶ However, the existence of sex-related differences in BP adaptation following IET has also been under-researched, with inconsistent interpretations of the findings to date. Following critical evaluation of the IET literature, pooled data from a meta-analysis by Inder *et al*⁷ and an individual participant data meta-analysis by Smart *et al*⁸ indicated no differences in the BP responses between males and females following IET. Conversely, a systematic review and narrative synthesis by Bentley *et al*⁹ demonstrated a greater IET effect for BP reductions in females, highlighting a lack of consensus regarding sex-related differences.

A greater understanding of sex-related specificity in exercise prescription may help further enhance the implementation of IET as a therapeutic intervention in the management of BP. While there is growing evidence to support the efficacy of IET in reducing BP in females, any differences in the magnitude of BP reduction between females and males and the mechanistic pathways responsible for BP adaptation remain unclear. Therefore, this study aimed to engage female and male participants with an established IET protocol compared with control, then (1) explore any differences in the BP adaptations between female and male participants following IET and (2) investigate any differences in the associated physiological mechanisms between sexes.

METHOD

Participants

According to the ESH guidelines, participants with normal to high-normal systolic BP (sBP) (range 120–140 mm Hg) were recruited. A current physical activity level below the WHO recommendations of ≥ 150 min of moderate physical activity per week was also required for participation, which included low-level physical activity, such as walking, light cycling and general household activities.¹⁰ Participants were free from injury or illness that may have affected the outcome variables of the study, were not taking any medication (or had a history of taking any anti-hypertensive medication), were non-smokers, consumed less than 14 units of alcohol per week and were not involved in purposeful exercise regimens. Screening was conducted using a standard Physical Activity Readiness Questionnaire, with an additional question added for females to assess whether they had regular menstruation, were taking oral contraception or were postmenopausal. At the time of testing, 12 female participants were taking hormonal contraception, 18 had regular menstruation

and 6 were postmenopausal. For females with regular menstruation (approximately a 28-day cycle), laboratory data collection was timed around their self-reported early follicular phase (days 1–7 of their menstrual cycle).

Study procedures

Recruitment was staggered on a rolling basis in batches of 10 to achieve the required sample size of 100 participants calculated using G*Power (Heinrich-Heine-Universität Düsseldorf, Düsseldorf, Germany).¹¹ For a direct comparison of the primary outcome variable (sBP) between the sexes, based on an alpha level of 0.05, a power of 0.80, two groups (female and male) and one covariate (baseline BP values), a sample size of 80 participants (40:40) was able to detect a small-to-moderate effect (0.32 Cohen's d). For the first phase of the study comparing all 80 intervention participants (female and male) to a control group of 20 participants, the estimated power based on likely changes observed by Taylor *et al*¹² of 12 mm Hg sBP and 6 mm Hg diastolic BP (dBP) was in excess of 95% for both sBP and dBP.

Following screening, participants were randomly allocated (at a ratio of 4:1 within each batch) to either the IET intervention (n=80) or control (n=20). Randomisation was carried out by the lead researcher using the RAND function on Microsoft Excel. The study set out to recruit a 50:50 split of females and males. Thus, within each batch, four female and four male participants were allocated to the intervention group, and one female and one male were assigned to the control group. All intervention participants completed a 4-week home-based IET programme. Participants were asked to attend the laboratory on four separate occasions. The first visit was a familiarisation session where participants were exposed to the laboratory and exercise equipment. During the second visit, participants were asked to undertake an incremental IET test to determine their individualised knee joint angle used during training, as previously described.¹³ As outlined below, the third and fourth visits recorded physiological data before and after the 4-week training period. Control participants were required to attend the laboratory twice for baseline and week-4 resting measures.

All participants were asked to maintain habitual dietary habits and daily routines throughout the study. Participants were asked to attend the laboratory at the same time (± 2 hours), fast for 4 hours, and avoid caffeine and alcohol for 24 hours before testing. For the IET group, post-training measures were taken between 72–96 hours after the participant's final training session to avoid the influence of acute hypotension that may persist postexercise.

Resting measures

All haemodynamic and cardiac autonomic variables were measured using a validated Task Force Monitor (TFM) (CNSystems, Medizintechnik, Graz, Austria).¹⁴ All measurements were taken in a temperature-controlled

room pre-IET and post-IET. To measure continuous BP, the TFM uses a vascular unloading technique at the proximal limb of the index or middle finger.¹⁵ Continuous measurement of BP is then adjusted in accordance with oscillometric BP values from the brachial artery of the contralateral arm.¹⁴ Heart rate (HR) was recorded using a six-channel ECG. Beat-to-beat stroke volume (SV) was measured with impedance cardiography via one electrode band applied to the nape of the neck and two placed on either side of the thorax, in line with the xiphisternum. Cardiac output (Q) was then subsequently calculated as the product of HR and SV. Total peripheral resistance, a proxy for the resistance to blood flow caused by systemic vasculature, was measured according to Ohm's law as a function of mean arterial pressure (MAP) and Q.

A six-channel ECG was used to measure R-R intervals, with the subsequent values used to calculate real-time HR variability (HRV) via an autoregressive model.¹⁶ The ECG traces were manually screened to clear any erroneous data. High frequency (HF) (predominantly parasympathetic outflow) and low frequency (LF) (predominantly sympathetic outflow)¹⁷ frequency parameters of HRV were automatically calculated by the TFM. These variables were expressed in absolute (ms^2) and normalised units (HFnu/LFnu). The sequence method automatically calculated baroreceptor reflex sensitivity (BRS)¹⁸.

Resting haemodynamic and cardiac autonomic measures were measured at baseline and 4 weeks. Following 15 min of seated rest, haemodynamic and cardiac autonomic function were measured continuously in the seated position for 5 min. After this time, the mean for each variable was calculated offline for 5 min.

IET intervention

The exercise training intervention consisted of an established lower-body IET programme¹³ thrice weekly over 4 weeks, with 48 hours between sessions. Each training session comprised four bouts of IET exercise separated by 2 min of seated rest.¹³ The training was performed at a participant-specific knee joint angle ascertained during an incremental IET test relative to 95% HR_{peak} (HR_{peak} relates to the mean HR evidenced during the final 30 s of the incremental test). Participants were required to record their HR during the last 30 s of each exercise bout using a wrist-mounted HR monitor and a Polar H10 chest strap (Polar Electro Oy, Kempele, Finland) and used to confirm adherence to training. If the HR data fell outside

the target HR zone for more than two sessions, the knee angle was reduced or increased by 5° to ensure that the HR fell back within the correct range.

Data analysis

Data were checked for normality assumptions. Where these were met, a one-way analysis of covariance (ANCOVA) was first carried out to explore the overall differences between all female and male participants ($n=80$) compared with a control group ($n=20$), with baseline measures as the covariate. Differences in each dependent variable (week-4 resting measures) between female ($n=37$) and male ($n=43$) participants were then explored using an ANCOVA with baseline measures as the covariate. Where data were not normally distributed, a Quade non-parametric test was used. An alpha level of <0.05 was set as the threshold for statistical significance.

Patient and public involvement

Participants were not involved in the study design, conduct or dissemination of outcomes.

RESULTS

113 participants were recruited to achieve a target sample size of 100 (table 1). The study experienced an 11.5% attrition rate, with 13 participants lost to follow-up. This included seven dropouts due to COVID-19-related reasons, three withdrawn participants for non-compliance with the protocol and three voluntarily withdrew. No adverse events were reported during the study. Of the 36 female participants allocated to the IET intervention, one missed a session (99.6% adherence). Two of the 44 male participants allocated to the IET intervention missed a single session each (99.8% adherence).

Overall BP-lowering effect compared with control

Following the 4-week training period, resting sBP ($M \pm SD$) decreased (-9.45 ± 6.69 mm Hg) and was significantly different to the control group (-0.79 ± 3.97 mm Hg, $p < 0.001$). Resting dBP reduced following IET (-5.62 ± 9.92 mm Hg) and significantly differed from the control group (2.19 ± 4.58 mm Hg, $p = 0.007$). Reductions in MAP (-6.90 ± 7.77 mm Hg) were also significantly different from the control group following IET (1.20 ± 3.84 mm Hg, $p < 0.001$) (figure 1).

Table 1 Baseline characteristics of all participants used for analysis

Variable	IET female (n=36)	IET male (n=44)	Control female (n=8)	Control male (n=12)
Age (years)	39 $\pm$ 12	36 $\pm$ 12	33 $\pm$ 5	32 $\pm$ 6
Height (cm)	166 $\pm$ 6	180 $\pm$ 8	167 $\pm$ 5	174 $\pm$ 8
Body mass (kg)	72 $\pm$ 9	88 $\pm$ 13	64 $\pm$ 7	74 $\pm$ 12
Data are presented as mean $\pm$ SD. Age, height and body mass were used for all intervention and control participants at baseline. IET, isometric exercise training.				




Figure 1 The mean systolic (a), diastolic (b) and mean arterial (c) pressure changes for the control (◆) and training (■) conditions. Error bars indicate the SE of the mean. * $p < 0.05$, ** $p < 0.01$ for the difference between experimental and control conditions.




Figure 2 Percentage distribution of BP categories in females (a) and males (b) pre-IET and post-IET. BP, blood pressure; IET, isometric exercise training.

BP variables (females vs males)

There were no significant differences in sBP ($p = 0.256$), dBP ($p = 0.119$) or MAP ($p = 0.066$) reductions between male and female participants following the 4-week training period. Despite no significant differences between the groups, it was found that 89% of females experienced a clinically important 5 mm Hg drop in sBP¹⁹ following IET compared with 76% of males. A 5 mm Hg reduction can be considered a clinically important difference as it reduces the risk of major cardiovascular events by about 10%, even among individuals with normal or high-normal BP values.¹⁹ Mean values for sBP, dBP and MAP can be found in online supplemental table 2.

BP reclassification

Before training, 50% of the female participants were categorised as having normal BP and 50% as having high-normal BP. After 4 weeks of IET, 50% had optimal ($<120/80$ mm Hg) BP, 44% had normal ($120\text{--}129/80\text{--}84$ mm Hg) BP and 6% had high-normal ($130\text{--}139/85\text{--}89$ mm Hg) BP. For male participants before training, 52% were categorised as having normal BP and 48% as having high-normal BP. After 4 weeks of IET, 38% had optimal BP, 45% had normal BP and 15% had high-normal BP (figure 2). The number needed to treat (NNT) to recategorise one female from normal to optimal BP is 2, and the NNT to recategorise one female from high-normal to normal BP is 3. For males, the NNT to recategorise one male from normal to optimal BP is 3 and the NNT to recategorise one male from high-normal to normal BP is 4.

Haemodynamic and cardiac autonomic variables (females vs males)

Following the 4-week training period, there were significant differences between the male and female participants in LF:HF ratio ($p=0.004$), LFnu ($p=0.015$) and HFnu ($p=0.015$) (online supplemental table 3). There were no significant differences in any of the other measured variables.

DISCUSSION

The primary aim of this study was to explore any differences in BP reduction and the associated underpinning physiological mechanisms between females and males following 4 weeks of IET. It was found that IET was effective in reducing resting BP compared with the control and that there were no significant differences in the magnitude of BP reduction between female and male participants. This is in agreement with previous meta-analytic findings by Inder *et al.*⁷ and Smart *et al.*⁸ However, to the best of our knowledge, this is the largest primary research study to date with a high representation of both females and males to specifically investigate the efficacy of IET for BP reduction. Previous studies have used sample sizes of <30 participants to compare sex-based differences^{7 8} compared with 80 in the current study. Therefore, these findings provide robust evidence for the utility of IET as an effective intervention for BP reduction in both sexes.

Despite no significant difference between females and males in BP reduction, the current study found statistically significant differences in specific HRV parameters. Following the 4-week training period, females were found to have lower LFnu and higher HFnu and a lower LF:HF ratio, which is known to represent sympatho-vagal balance²⁰ compared with male participants. These responses indicate increased cardiac vagal tone and parasympathetic modulation.²¹ It has been documented that lower-body IET, specifically the IET protocol used in the present study, enhances HRV and sympathovagal balance in males.¹² However, the greater autonomic response observed in females in the current investigation may be linked to sex-related differences in hormonal and vascular biology that are known to influence cardiovascular regulation.²² Females typically show greater parasympathetic activity affecting the heart and have less sympathetic input in vascular tone.²³ Female sex hormones, specifically oestrogen and progesterone, are known to improve HRV and vascular function, respectively. Together, these hormones provide vascular protective effects that are mediated through a complex interplay of physiological mechanisms, such as modulation of L-type calcium channel activity,²⁴ upregulation of the nitric oxide (NO) pathway^{25 26} and greater expression of neuronal NO synthase in vagal neurons.²⁷ Moreover, oestrogen has been shown to affect higher brain centres, including the hypothalamus and amygdala, which regulate autonomic functions and enhance parasympathetic activity.²⁸ Furthermore, it has been shown that beta-adrenergic

receptors are influenced by oestrogen, which can lead to an overall reduction in sympathetic drive.^{29 30} Although oestrogen and progesterone levels were not measured in the current study, it is possible that the cardioprotective effects from these female sex hormones augmented the autonomic adaptations specifically seen in the female participants following the 4-week IET period. This is an important finding as increased HRV and vagal function are associated with a decreased risk of morbidity and mortality, independent of traditional risk factors.³¹

Furthermore, studies that have compared sex-based differences following acute bouts of exercise have also reported greater parasympathetic activity in females. Kappus *et al.*³² found greater vagal reactivation in females during recovery from maximal aerobic exercise. Teixeira *et al.*³³ also found a greater increase in cardiac BRS (used as a measure of reflexive vagal modulation) following an acute bout of isometric handgrip training. Arguably, these findings are expected since strong associations have previously been observed between acute and chronic responses during exercise, which has led to the suggestion that chronic adaptations may result from the temporal summation of acute responses.³⁴

Clinical and cost effectiveness of IET delivery

Although there were no significant differences in the BP reductions following the 4 weeks, it should be acknowledged that 89% of females experienced a clinically important 5 mm Hg drop in sBP compared with 76% of males. A 5 mm Hg reduction in sBP reduces the risk of major cardiovascular events by approximately 10%, even among individuals with normal or high-normal BP values.¹⁹ Moreover, more female participants were reclassified into the optimal BP category following IET (figure 2), which may be related to further reductions in cardiovascular risk.³⁵ The higher incidence of a clinically important reduction in BP and greater reclassification into the optimal BP category may be related to the autonomic adaptations evidenced and could indicate a greater cardioprotective effect for females following IET.

We also found that the NNT to recategorise one female from normal to optimal BP is 2, and the NNT to recategorise one female from high-normal to normal BP is 3. For males, we found the NNT to recategorise one male from normal to optimal BP is 3, and the NNT to recategorise one male from high-normal to normal BP is 4.³⁶ Related to this, our mean reduction in sBP of 10.5 mm Hg for females and 8.8 mm Hg for males exceeds the 7 mm Hg reported in the meta-analysis by Xie *et al.*³⁶ These authors demonstrated that a reduction of 7 mm Hg in sBP equated to a 13% risk reduction for myocardial infarction and a 22% risk reduction for stroke. Intuitively, we could expect similar or even greater reductions in these events in women and men with HTN who accomplish a reduction in sBP of 7 mm Hg or more from undertaking IET. Of course, these benefits should be considered regarding treatment delivery costs, especially as the expected sBP reduction from IET is similar

to antihypertensive monotherapy.³⁷ The cost of a single antihypertensive medication to healthcare services in the UK can be up to £370 for direct renin inhibitors.³⁸ Based on a direct comparison of end-user products only (ie, home-based equipment used for IET), the IET intervention would have an estimated cost of £10 per annum per patient.³⁹

Limitations

We did not measure oestrogen levels in participants, and therefore, inferences based on its effect on the cardiovascular system lack physiological verification. Moreover, the type of contraceptive pill used was not controlled in this study. Although previous research has shown no differences in autonomic modulation throughout a full cycle while using oral contraceptive pills,⁴⁰ there are data to suggest fluctuations in endogenous and exogenous hormone levels throughout the pill cycle, which may have affected HRV.⁴¹ Results from the current study are also only applicable to a 4-week training period, and it remains to be established whether there are sex differences with a more prolonged training dose.

CONCLUSIONS

Our findings support the efficacy of lower-body IET in equally reducing BP in females and males. However, the greater autonomic response seen in females and a higher prevalence of clinically significant BP reductions may indicate a greater cardioprotective effect. These findings may have important implications for future IET prescription and could help practitioners develop sex-specific protocols to optimise cardiovascular outcomes.

Author affiliations

¹School of Sport, Health and Applied Science, St Mary's University Twickenham, Twickenham, UK

²Sport, Exercise, and Rehabilitation Sciences, Canterbury Christ Church University, Canterbury, Kent, UK

³Institute of Sport, Exercise and Health, Division of Surgery and Interventional Science, University College London, London, UK

⁴School of Science and Technology, University of New England, Armidale, New South Wales, Australia

⁵University of Kent Centre for Health Services Studies, Canterbury, UK

Contributors HS and JW completed the acquisition of data. HS, JW and DC performed the data analysis. HS and JW were the principal writers of the manuscript. All authors contributed to the drafting and revision of the final article. All authors approved the final submitted version of the manuscript. JW is the guarantor.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants. The study was approved by the ethics committee of Canterbury Christ Church University (Ref: 19/SAS/12C). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Harry Swift <http://orcid.org/0000-0003-0499-5706>

Damian Coleman <http://orcid.org/0000-0003-1861-322X>

Charles Pedlar <http://orcid.org/0000-0002-3075-9101>

Neil Andrew Smart <http://orcid.org/0000-0002-8290-6409>

Chris Farmer <http://orcid.org/0000-0003-1736-8242>

Jonathan Wiles <http://orcid.org/0000-0002-7790-8063>

REFERENCES

- Roth GA, Abate D, Abate KH, *et al*. Erratum: Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980–2017: a systematic analysis for the Global Burden of Disease Study 2017 (The Lancet [2018] 392(10159) (1736–1788)[S0140673618322037](10.1016/S0140-6736(18)32203-7). *Lancet* 2018;392.
- Saheera S, Krishnamurthy P. Cardiovascular Changes Associated with Hypertensive Heart Disease and Aging. *Cell Transplant* 2020;29:963689720920830.
- Zhou B, Carrillo-Larco RM, Danaei G. Worldwide trends in hypertension prevalence and progress in treatment and control from 1990 to 2019: a pooled analysis of 1201 population-representative studies with 104 million participants. *Lancet* 2021;398:957–80.
- Mancia G, Kreutz R, Brunström M, *et al*. 2023 ESH Guidelines for the management of arterial hypertension The Task Force for the management of arterial hypertension of the European Society of Hypertension: Endorsed by the International Society of Hypertension (ISH) and the European Renal Association (ERA). *J Hypertens* 2023;41:1874–2071.
- Cornelissen VA, Smart NA. Exercise training for blood pressure: a systematic review and meta-analysis. *J Am Heart Assoc* 2013;2:e004473.
- Edwards JJ, Deenmamode AHP, Griffiths M, *et al*. Exercise training and resting blood pressure: a large-scale pairwise and network meta-analysis of randomised controlled trials. *Br J Sports Med* 2023;57:1317–26.
- Inder JD, Carlson DJ, Dieberg G, *et al*. Isometric exercise training for blood pressure management: a systematic review and meta-analysis to optimize benefit. *Hypertens Res* 2016;39:88–94.
- Smart NA, Way D, Carlson D, *et al*. Effects of isometric resistance training on resting blood pressure: individual participant data meta-analysis. *J Hypertens* 2019;37:1927–38.
- Bentley DC, Nguyen CH, Thomas SG. Resting blood pressure reductions following handgrip exercise training and the impact of age and sex: a systematic review and narrative synthesis. *Syst Rev* 2018;7:229.
- World Health Organization. Global action plan on physical activity 2018–2030. 2019.
- Faul F, Erdfelder E, Buchner A, *et al*. Statistical power analyses using G*Power 3.1: tests for correlation and regression analyses. *Behav Res Methods* 2009;41:1149–60.
- Taylor KA, Wiles JD, Coleman DA, *et al*. Neurohumoral and ambulatory haemodynamic adaptations following isometric exercise training in unmedicated hypertensive patients. *J Hypertens* 2019;37:827–36.
- Wiles JD, Goldring N, Coleman D. Home-based isometric exercise training induced reductions resting blood pressure. *Eur J Appl Physiol* 2017;117:83–93.
- Fortin J, Habenbacher W, Gruellenberger R, *et al*. Real-time monitor for hemodynamic beat-to-beat parameters and power spectra analysis of the biosignals. 2002.

- 15 Parati G, Ongaro G, Bilo G, *et al.* Non-invasive beat-to-beat blood pressure monitoring: new developments. *Blood Press Monit* 2003;8:31–6.
- 16 Fortin J, Haïtchi G, Bojic A, *et al.* Validation and verification of the task force® monitor. FDA paper; 2001.
- 17 Akselrod S, Gordon D, Ubel FA, *et al.* Power spectrum analysis of heart rate fluctuation: a quantitative probe of beat-to-beat cardiovascular control. *Science* 1981;213:220–2.
- 18 Valipour A, Schneider F, Kössler W, *et al.* Heart rate variability and spontaneous baroreflex sequences in supine healthy volunteers subjected to nasal positive airway pressure. *J Appl Physiol (1985)* 2005;99:2137–43.
- 19 Canoy D, Nazarzadeh M, Copland E, *et al.* How Much Lowering of Blood Pressure Is Required to Prevent Cardiovascular Disease in Patients With and Without Previous Cardiovascular Disease? *Curr Cardiol Rep* 2022;24:851–60.
- 20 Pagani M, Montano N, Porta A, *et al.* Relationship Between Spectral Components of Cardiovascular Variabilities and Direct Measures of Muscle Sympathetic Nerve Activity in Humans. *Circulation* 1997;95:1441–8.
- 21 Kim H-G, Cheon E-J, Bai D-S, *et al.* Stress and Heart Rate Variability: A Meta-Analysis and Review of the Literature. *Psychiatry Investig* 2018;15:235–45.
- 22 Gerdtts E, Sudano I, Brouwers S, *et al.* Sex differences in arterial hypertension. *Eur Heart J* 2022;43:4777–88.
- 23 Fisher JP, Adlan AM, Shantsila A, *et al.* Muscle metaboreflex and autonomic regulation of heart rate in humans. *J Physiol (Lond)* 2013;591:3777–88.
- 24 Barbagallo M, Dominguez LJ, Licata G, *et al.* Vascular Effects of Progesterone : Role of Cellular Calcium Regulation. *Hypertension* 2001;37:142–7.
- 25 Aryan L, Younessi D, Zargari M, *et al.* The Role of Estrogen Receptors in Cardiovascular Disease. *Int J Mol Sci* 2020;21:4314.
- 26 Khalil RA. Sex hormones as potential modulators of vascular function in hypertension. *Hypertension* 2005;46:249–54.
- 27 Xia Y, Krukoff TL. Estrogen induces nitric oxide production via activation of constitutive nitric oxide synthases in human neuroblastoma cells. *Endocrinology* 2004;145:4550–7.
- 28 McEwen BS, Akama KT, Spencer-Segal JL, *et al.* Estrogen effects on the brain: actions beyond the hypothalamus via novel mechanisms. *Behav Neurosci* 2012;126:4–16.
- 29 Machuki JO, Zhang HY, Harding SE, *et al.* Molecular pathways of oestrogen receptors and β -adrenergic receptors in cardiac cells: Recognition of their similarities, interactions and therapeutic value. *Acta Physiol (Oxf)* 2018;222:e12978.
- 30 Hart EC, Charkoudian N, Wallin BG, *et al.* Sex and ageing differences in resting arterial pressure regulation: the role of the β -adrenergic receptors. *J Physiol (Lond)* 2011;589:5285–97.
- 31 Thayer JF, Lane RD. The role of vagal function in the risk for cardiovascular disease and mortality. *Biol Psychol* 2007;74:224–42.
- 32 Kappus RM, Ranadive SM, Yan H, *et al.* Sex differences in autonomic function following maximal exercise. *Biol Sex Differ* 2015;6:28.
- 33 Teixeira AL, Ritti-Dias R, Antonino D, *et al.* Sex Differences in Cardiac Baroreflex Sensitivity after Isometric Handgrip Exercise. *Med Sci Sports Exerc* 2018;50:770–7.
- 34 Farah BQ, Germano-Soares AH, Rodrigues SLC, *et al.* Acute and Chronic Effects of Isometric Handgrip Exercise on Cardiovascular Variables in Hypertensive Patients: A Systematic Review. *Sports (Basel)* 2017;5:55.
- 35 Wright JT Jr, Whelton PK, Johnson KC, *et al.* SPRINT Revisited: Updated Results and Implications. *Hypertension* 2021;78:1701–10.
- 36 Xie X, Atkins E, Lv J, *et al.* Effects of intensive blood pressure lowering on cardiovascular and renal outcomes: updated systematic review and meta-analysis. *Lancet* 2016;387:435–43.
- 37 Wong GWK, Boyda HN, Wright JM. Blood pressure lowering efficacy of beta-1 selective beta blockers for primary hypertension. *Cochrane Database Syst Rev* 2016;3:CD007451.
- 38 NICE. Hypertension in adults: diagnosis and management. *Practice Nurse* 2019;49.
- 39 Lea JWD, O'Driscoll JM, Wiles JD. The implementation of a home-based isometric wall squat intervention using ratings of perceived exertion to select and control exercise intensity: a pilot study in normotensive and pre-hypertensive adults. *Eur J Appl Physiol* 2024;124:281–93.
- 40 Teixeira AL, Ramos PS, Vianna LC, *et al.* Heart rate variability across the menstrual cycle in young women taking oral contraceptives. *Psychophysiology* 2015;52:1451–5.
- 41 Brahmabhatt Y, Gupta M, Hamrahian S. Hypertension in Premenopausal and Postmenopausal Women. *Curr Hypertens Rep* 2019;21:74.